

A COINCIDENCE BETWEEN THE GENERAL EFFICIENCY AND THE APPROXIMATE DOMINANCE

Vasile Postolică

Department of Mathematics, Informatics and Educational Sciences,

”Vasile Alecsandri” University of Bacău, Romania

vpostolica@ambra.ro

Abstract This research work represents a short synthesis on the main results obtained until this moment for the general Efficiency and its immediate connections with the Approximate Dominance expressed by the Convex Cones, in the ample context of the Ordered Vector Spaces. The largest class of such as these Ordering Cones, named by us Isac’s Cones in Hausdorff Locally Convex Spaces ensures these maximum links, together with the proved applications including the Choquet Boundaries.

**Work presented as invited lecture at CAIM 2014, September 19-22,
”Vasile Alecsandri” University of Bacău, Romania.**

Keywords: general efficiency, approximate dominance, Isac’s (nuclear, supernormal) cone, Choquet boundary.

2010 MSC: 90B50, 90B70.

1. INTRODUCTION

Until now, the basis on which was developed our general concept of the Efficiency in the Ordered Linear Spaces and its Applications, supplied by the Ordered Locally Convex Spaces, is represented about the vastness class of the Convex Cones introduced by Professor Isac in 1981, published in 1983, called by us and, officially recognized, as “Isac’s Cone” in 2009, after the acceptance of this last agreed denomination by Professor Isac. To it is devoted this scientific research work, following the content of this International Meeting. All the elements concerning the ordered topological vector spaces used here are in accordance with *Nachbin, L., 1965* and *Peressini, A., L., 1967*.

2. THE RECENT EFFICIENCY AND THE APPROXIMATE DOMINANCE IN THE ORDERED VECTOR SPACES

Let E be any real or complex vector space ordered by a convex cone K , K_1 a non-void subset of K and A a non-empty subset of E . The following definition introduces a

new concept of the approximate efficiency, which generalizes the well known notion of Pareto minimum type efficiency in every Euclidean space and not only.

Definition 2.1. (Postolică, V., 2002, 2008). We say that $a_0 \in A$ is a K_1 -minimal efficient point of A , in notation, $a_0 \in MIN(A, K, K_1)$ if it satisfies one of the following equivalent conditions:

- (i) $A \cap (a_0 - K - K_1) \subseteq a_0 + K + K_1$;
- (ii) $(K + K_1) \cap (a_0 - A) \subseteq -K - K_1$.

In a similar manner one defines the K_1 -maximal efficient points by replacing $K + K_1$ with $-(K + K_1)$. The set of such points will be denoted by $MAX(A, K, K_1)$. Whenever $K_1 = \{0\}$ we consider $MIN(A, K, K_1) = MIN(A, K)$ and $MAX(A, K, K_1) = MAX(A, K)$, respectively.

All the efficient points of the set A with respect to the convex cone K following K_1 are represented by $eff(A, K, K_1) = MIN(A, K, K_1) \cup MAX(A, K, K_1)$. Consequently, $eff(A, K) = MIN(A, K) \cup MAX(A, K)$. Clearly,

$$\begin{aligned} A \cap (a_0 \mp K) \subseteq a_0 \pm K_1 &\Rightarrow A \cap (a_0 \pm K \pm K_1) \subseteq a_0 \mp K \mp K_1 \Rightarrow \\ &\Rightarrow A \cap (a_0 \mp K_1) \subseteq a_0 \pm K, \end{aligned}$$

which suggests other concepts for the efficiency and its approximate term in the general ordered linear spaces.

Remark 2.1. Obviously, $a_0 \in eff(A, K, K_1)$ iff it is a fixed point for at least one of the multifunctions $F_{\mp} : A \rightarrow 2^A$ defined by

$$F_{\mp}(t) = \left\{ a \in A : A \cap (a \mp K \mp K_1) \subseteq t \pm K \pm K_1 \right\},$$

with the corresponding signs. Consequently, for the existence of these general efficient points we can also apply appropriate fixed points theorems concerning the multi-functions (see, for instance, Cardinali, T., Papalini, F., 1994, Park, S., 1992, Patriche, M., 2013, Zhang Cong – Jun, 2005 and any other proper scientific papers).

Remark 2.2. Németh, A.B., 1989, proved that, whenever $K_1 \subseteq K \setminus \{0\}$, the existence of this new type of the efficient points for the lower bounded sets characterizes the semi-Archimedean ordered vector spaces and the regular ordered locally convex spaces.

Remark 2.3. Whenever K is pointed, that is, $K \cap (-K) = \{0\}$, then $a_0 \in MIN(A, K, K_1)$ ($a_0 \in MAX(A, K, K_1)$) means that $A \cap (a_0 \mp K \mp K_1) = \emptyset$ or, equivalently, $(\pm K + K_1) \cap (a_0 - A) = \emptyset$ for $0 \notin K_1$ and $A \cap (a_0 \mp K \mp K_1) = \{a_0\}$, respectively, if $0 \in K_1$. Whenever $K_1 = \{0\}$, from Definition 1 one obtains the usual concept of the efficient (Pareto minimal, optimal or admissible) point : $a_0 \in MIN(A, K)(MAX(A, K))$ if it fulfils

- (i), (ii) or any of the next equivalent properties:
- (iii) $(A \pm K) \cap (a_0 \mp K) \subseteq a_0 \pm K$;
- (iv) $K \cap (a_0 - A \mp K) \subseteq \mp K$.

This shows that a_0 is a fixed point for at least one of the following multifunctions:

$$F_1 : A \rightarrow A, F_1(t) = \left\{ a \in A : A \cap (a \mp K) \subseteq t \pm K \right\},$$

$$F_2 : A \rightarrow A, F_2(t) = \left\{ a \in A : A \cap (t \mp K) \subseteq a \pm K \right\},$$

$$F_3 : A \rightarrow A, F_3(t) = \left\{ a \in A : (A \pm K) \cap (a \mp K) \subseteq t \pm K \right\},$$

$$F_4 : A \rightarrow A, F_4(t) = \left\{ a \in A : (A \pm K) \cap (t \mp K) \subseteq a \pm K \right\},$$

that is, $a_0 \in F_i(a_0)$ for some $i = \overline{1,4}$. If, in addition, K is pointed, then $a_0 \in A$ is an efficient point of A with respect to K if and only if one of the following equivalent relations hold :

- (v) $A \cap (a_0 \mp K) = \{a_0\}$;
- (vi) $A \cap (a_0 \mp K \setminus \{0\}) = \emptyset$;
- (vii) $(\pm K) \cap (a_0 - A) = \{0\}$;
- (viii) $(\pm K \setminus \{0\}) \cap (a_0 - A) = \emptyset$;
- (ix) $(A \pm K) \cap [a_0 \mp (K \setminus \{0\})] = \emptyset$.

We also notice that

$$MIN(A, K) (MAX(A, K)) = \bigcap_{\{0\} \neq K_2 \subseteq K} MIN(A, K, K_2) \left(\bigcap_{\{0\} \neq K_2 \subseteq -K} MAX(A, K, K_2) \right).$$

Moreover, $a_0 \in \text{eff}(A, K)$ if only if it is a critical (ideal or balance) point (see, for example, Kim, W. K., Tan, K.K., 2001; Isac, G. 1985; Isac, G., Bulavsky, V. A., Kalashnikov, V., V. 2002; Isac, G., Postolică, V., 1993; Postolică, V., Scarelli, A., Venzi, L., 2001; Postolică, V., 2004 and the corresponding references) for the generalized dynamical systems $\Gamma_{\mp} : A \rightarrow 2^A$ defined by $\Gamma_{\mp}(a) = A \cap (a \mp K)$, $a \in A$. In this way, $\text{eff}(A, K)$ describes the balance extremum moments for Γ_{\mp} , which in the market context, expresses the competitive equilibrium/non-equilibrium consisting of the general relation price/consumption.

Considering $K_1 = \{\varepsilon\}(\varepsilon \in K \setminus \{0\})$, one obtains that $a_0 \in \text{eff}(A, K, K_1)$ iff $A \cap (a_0 - \varepsilon \mp K) = \emptyset$. In all these cases, for the set $\text{MIN}(A, K, K_1)$ we used the notation $\varepsilon - \text{MIN}(A, K)$. It is obvious that $\text{MIN}(A, K) = \bigcap_{\varepsilon \in K \setminus \{0\}} [\varepsilon - \text{MIN}(A, K)]$, with the natural projections for $\text{MAX}(A, K)$ obtained by replacing the convex cone K with $-K$. When referring to the existence of the efficient points, the significant properties of the sets containing these points and the applications in the usual or extended contexts we mention, as a comprehensive bibliography, (Aubin, J.P., 1993; Benson, H. P., 1977, 1979, 1983; Borwein, J. M., 1977, 1983; Dauer, J. P., Gallagher, R. J., 1990; Dominiak, Cezary, 2006; Ehrgott, Matthias, 2005; Fishburn, P. C., 1984; Gass, Saul, I., Harris, Carl, M., 2001; Geoffrion, A. M., 1968; Hartley, R., 1978; Henig, M. I., 1982; Heyne, Paul, 2000; Hyers, D., H., Isac, G., Rassias, T.M., 1997; Isac, G., 1981, 1983, 1985, 1994, 1998, 2003, Isac, G., Bahya, A. O., 2002; Isac, G., Postolică, V., 1993, 2005; Isac, G., Tammer, Chr., 2003; Jablonsky, J., 2006; Loricidan, P., 1984; Luc, D. T., 1989; Luptăci, M., Böhm, B., 2006; Németh, A., B., 1989; Ng, K., F., Zheng, X., Y., 2002; Postolică, V., 1989, 1993, 1994, 1997, 1999, 2001, 2002, 2004, 2008, 2009; Songsak Sriboonchitta, Wing – Keung Wong, Sompong Dhopongsa, Hung, T., Nguyen, 2009; Sontag, Z., ZĂlinescu, C., 2000; Stewart, D., T., 2004; Truong, X. D. H., 1994; Zimmermann, H., 2000) with widely established results.

Let us consider the set $S \text{MIN}(A, K, K_1) = \{a_1 \in A : A \subseteq a_1 + K + K_1\}$ of all *strong minimum points* of A with respect to K by K_1 and $S \text{MAX}(A, K, K_1) = \{a_2 \in A : A \subseteq a_2 - K - K_1\}$ the set of all *strong maximum points* of A with respect to K by K_1 , respectively, and $S(A, K, K_1) = \{a_1 \in A : A \subseteq a_1 + K + K_1\} \cup \{a_2 \in A : A \subseteq a_2 - K - K_1\}$

The following theorem shows the immediate connection between the approximate efficiency and the strong optimization through the agency of cone minimal (cone maximal) efficient points.

Theorem 2.1. *If $S(A, K, K_1) \neq \emptyset$, then $S(A, K, K_1) = \text{eff}(A, K, K_1)$.*

Proof. In fact, we prove that

$$S \text{MIN}(A, K, K_1) \neq \emptyset \Rightarrow \text{MIN}(A, K, K_1) = S \text{MIN}(A, K, K_1)$$

and

$$S \text{MAX}(A, K, K_1) \neq \emptyset \Rightarrow \text{MAX}(A, K, K_1) = S \text{MAX}(A, K, K_1),$$

so it is sufficient to show the first implication.

Clearly, $S \text{MIN}(A, K, K_1) \subseteq \text{MIN}(A, K, K_1)$. Indeed, if $a_0 \in S \text{MIN}(A, K, K_1)$ and $a \in A \cap (a_0 - K - K_1)$ are arbitrary elements, then $a \in a_0 + K + K_1$, that is, $a_0 \in \text{MIN}(A, K, K_1)$ by virtue of the point (i) contained in the Definition 2.1. Suppose now that $\bar{a} \in S \text{MIN}(A, K, K_1) \neq \emptyset$ and there exists $a_0 \in \text{MIN}(A, K, K_1) \setminus S \text{MIN}(A, K, K_1)$. Out of the fact $\bar{a} \in S \text{MIN}(A, K, K_1)$, it follows that $a_0 \in \bar{a} + K + K_1$, that is, $\bar{a} \in a_0 - K - K_1$, from which, since $\bar{a} \in A$ and $a_0 \in \text{MIN}(A, K, K_1)$, we conclude that $\bar{a} \in a_0 + K + K_1$. Therefore, $A \subseteq \bar{a} + K + K_1 \subseteq a_0 + K + K_1$, in contradiction with $a_0 \notin S \text{MIN}(A, K, K_1)$, as claimed. ■

Remark 2.4. It is obvious that $S(A, K, K_1) \subseteq \text{eff}(A, K, K_1)$. The previous theorem shows that every time there exists at least one strong cone minimum or cone maximum point, the set of all efficient points coincides with the set of all these points, respectively. Using this conclusion and the abstract construction given by us in (*Postolică, V. 1991, 1988*) for the splines in the H-locally convex spaces introduced in (*Precupanu, T., 1969*) as separated locally convex spaces with any semi-norm satisfying the parallelogram law, linear topological spaces also studied with strictness in (*Kramar, E., 1981*), it follows that the only best simultaneous and vectorial approximation for each element in the direct sum of any (closed) linear subspace and its orthogonal, with respect to a linear (continuous) operator between two arbitrary H-locally convex spaces, is its spline function. We also note that it is possible to have $S(A, K, K_1) = \emptyset$ and $\text{eff}(A, K, K_1) = A$.

Thus, for example, if one considers $X = R^n (n \in N, n \geq 2)$, $K = R_+^n$, $K_1 = \{(0, \dots, 0)\}$ and for each real number c we define

$$A_{0c} = \left\{ (x_i) \in X : \sum_{i=1}^n x_i = c \right\}, \quad A_{1c} = \left\{ (x_i) \in X : \sum_{i=1}^n x_i \geq c \right\},$$

$$A_{2c} = \left\{ (x_i) \in X : \sum_{i=1}^n x_i \leq c \right\},$$

then it is clear that

$$S(A_{0c}, K, K_1) = S(A_{1c}, K, K_1) = S(A_{2c}, K, K_1) = \emptyset,$$

$$\text{eff}(A_{0c}, K, K_1) = \text{MIN}(A_{0c}, K, K_1) = \text{MAX}(A_{0c}, K, K_1) = A_{0c},$$

$$\text{MIN}(A_{1c}, K, K_1) = \text{MAX}(A_{2c}, K, K_1) = A_{0c}$$

so, by extrapolation,

$$\text{eff}(A_{1c}, K, K_1) = A_{0c} \cup \{(+\infty, +\infty, \dots, +\infty)\}, \quad \text{eff}(A_{2c}, K, K_1) = A_{0c} \cup \{(-\infty, -\infty, \dots, -\infty)\}.$$

Meanwhile, in the usual real linear space of all sequences, ordered by the convex cone

$K = \{(x_n) : n \in N^*, n \geq 2, x_n \geq 0, \forall n \geq 2\}$, for $A = \{(x_{n\alpha}) : n \in N^*, n \geq 2, \alpha > 0\}$ with $x_{n\alpha} = (n-1)^{-\alpha} - n^{-\alpha}, n \in N^*, n \geq 2, \alpha > 0$ and $K_1 = \{(0, 0, \dots)\}$, we have $\text{eff}(A, K, K_1) = S(A, K, K_1) = \emptyset$.

Simple examples show that, in general, the implication

$$S(A, K, K_1) = \emptyset \Rightarrow \text{eff}(A, K, K_1) = A$$

is not true.

3. ISAC'S CONES AND THE EFFICIENCY

Initially, the next concept of convex cone named by its author *nuclear, super-normal* (Isac, G., 1980, 1981, 1983, 1985) and, by us, *Isac's cone* (Postolică, V., 2009) was introduced in the following context, being the most appropriate for the general efficiency by optimization and conversely (see, for instance, Isac, G., 1980, 1981, 1983, 1985, 1994, 1998, 2003; Isac, G., Bahya, A.O., 2002; Isac, G., Postolică, V., 1993, 2005; Isac, G., Tammer, Chr., 2003). As we will see, to ensure the existence of the efficient points for a non-empty set in Hausdorff separated locally convex space, the main reason will be to replace *the compactness* with *the completeness* hypothesis imposed on the set and the ordering cone to be an Isac's cone.

Definition 3.1. (Treves, F., 1967). A locally convex space is any couple $(X, \text{Spec}(X))$ which is composed of a real linear space X and a family $\text{Spec}(X)$ of seminorms on X such that:

- (i) $\chi p \in \text{Spec}(X), \forall \chi \in \mathbb{R}_+, p \in \text{Spec}(X)$;
- (ii) if $p \in \text{Spec}(X)$ and q is an arbitrary seminorm on X such that $q \leq p$, then $q \in \text{Spec}(X)$;
- (iii) $\sup(p_1, p_2) \in \text{Spec}(X), \forall p_1, p_2 \in \text{Spec}(X)$ where $\sup(p_1, p_2)(x) = \sup(p_1(x), p_2(x)), \forall x \in X$.

It is well known that whenever such a family as this $\text{Spec}(X)$ is given on a real vector space X , there exists a locally convex topology τ on X such that (X, τ) is a topological linear space and a seminorm p on X is τ -continuous iff $p \in \text{Spec}(X)$.

A non-empty subset B of $\text{Spec}(X)$ is a *base* for it if for every $p \in \text{Spec}(X)$ there exist $\chi > 0$ and $q \in B$ such that $p \leq \chi q$ and (X, τ) is a Hausdorff locally convex space iff $\text{Spec}(X)$ has a base B , named *Hausdorff base*, with the property that $\{x \in X : p(x) = 0, \forall p \in B\} = \{\theta\}$ where θ is the null vector in X .

In this research paper we will suppose that the space (X, τ) sometimes denoted by X is a Hausdorff locally convex space. Every non-empty subset K of X satisfying the following properties: $K + K \subseteq K$ and $\chi K \subseteq K, \forall \chi \in \mathbb{R}_+$ is named *convex cone*. If, in addition, $K \cap K = \{\theta\}$, then K is called *pointed*. Clearly, any pointed convex cone K in X generates an ordering on X defined by $x \leq y, y \in X$ iff $y - x \in K$. If X^* is the dual of X , then the *dual cone* of K is defined by $K^* = \{x^* \in X^* : x^*(x) \geq 0, \forall x \in K\}$ and its corresponding *polar* is $K^0 = -K^*$. We recall that a pointed convex cone $K \subset (X, \text{Spec}(X))$ is *normal* with respect to the topology defined by $\text{Spec}(X)$ if it fulfils one of the next equivalent assertions:

- (i) there exists a base Ω of neighborhoods for the origin θ in X such that

$$V = (V + K) \cap (V - K), \forall V \in \Omega;$$

- (ii) there exists a base B of $\text{Spec}(X)$ with $p(x) \leq p(y), \forall x, y \in K, x \leq y, \forall p \in B$;
- (iii) for any two nets $\{x_i\}_{i \in I}, \{y_i\}_{i \in I} \subset K$ with $\theta \leq x_i \leq y_i, \forall i \in I$ and $\lim_{i \in I} y_i = \theta$ it follows that $\lim_{i \in I} x_i = \theta$. In particular, a convex cone K is normal in a normed linear

space $(E, \|\cdot\|)$ iff there exists $t \in (0, \infty)$ such that $x, y \in E$ and $y - x \in K$ implies that $\|x\| \leq t \|y\|$.

It is well known (Treves, F., 1967) that, the concept of *normal cone* together with its natural generalizations represents the most important notion in the theory and the applications of the convex cones in the general topological ordered vector spaces. Thus, for example, for every separated locally convex space $(X, \text{Spec}(X))$ and any closed normal cone $K \subset (X, \text{Spec}(X))$ we have $X^* = K^* - K^*$ (for more details see Hyers, D., H., Isac, G., Rassias, T.M., 1997; Isac, G., 1980).

Each pointed convex cone $K \subset (X, \text{Spec}(X))$ for which there exists a non-empty, convex bounded set $B \subset X$ such that $0 \notin \bar{B}$ and $K = \bigcup_{\chi \geq 0} \chi B$ is called *well-based*.

A cone $K \subset (X, \text{Spec}(X))$ is *well-based* iff there exists a base $B = \{p_i\}_{i \in I}$ of $\text{Spec}(X)$ and a linear continuous functional $f \in K^*$ such that for every $p_i \in B$ there exists $c_i > 0$ with $c_i p_i(x) \leq f(x), \forall x \in K$ (Hyers, D., H., Isac, G., Rassias, T.M., 1997; Isac, G., 1980).

Clearly, every well - based cone is a normal cone, but, in general, the converse is not true, as we can see in the examples below, starting from the next basic notion.

Definition 3.2. (Isac, G., 1981, 1983). In a Hausdorff locally convex space $(X, \text{Spec}(X))$ a pointed convex cone $K \subset X$ is *nuclear (supernormal)* with respect to the topology induced by $\text{Spec}(X)$ if there exists a base $B = \{p_i\}_{i \in I}$ of $\text{Spec}(X)$ such that for every $p_i \in B$ there exists $f_i \in X^*$ with $p_i(x) \leq f_i(x), \forall x \in K$.

Remark 3.1. For the first time, we called any such as this cone “*Isac’s cone*” in (Postolici, V., 2009), taking into account that the above definition of any locally convex space is equivalent with the following: let X be a real or complex linear space and $P = \{p_\alpha : \alpha \in A\}$ a family of seminorms defined on X . For every $x \in X, \varepsilon > 0$ and $n \in \mathbb{N}^*$ let

$$V(x; p_1, p_2, \dots, p_n; \varepsilon) = \{y \in X : p_\alpha(y - x) < \varepsilon, \forall \alpha = \overline{1, n}\},$$

$$V_0(x) = \left\{ V(x; p_1, p_2, \dots, p_n; \varepsilon) : n \in \mathbb{N}^*, p_\alpha \in P, \alpha = \overline{1, n}, \varepsilon > 0 \right\}$$

has the properties :

$$(V1) \quad x \in V, \forall V \in V_0(x) ;$$

$$(V2) \quad \forall V_1, V_2 \in V_0(x), \exists V_3 \in V_0(x) : V_3 \subseteq V_1 \cap V_2 ;$$

$$(V3) \quad \forall V \in V_0(x), \exists U \in V_0(x), U \subseteq V \text{ such that } \forall y \in U, \exists W \in V_0(y) \text{ with } W \subseteq V.$$

Therefore, $V_0(x)$ is a *base of neighborhoods* for x and taking $\nu(x) = \{V \subseteq X : \exists U \in V_0(x) \text{ with } U \subseteq V\}$, the set $\tau = \{D \subseteq X : D \in \nu(x), \forall x \in D \cup \{\emptyset\}\}$ is the locally convex topology generated by the family P . Obviously, the usual operations which induce the structure of linear space on X are continuous with respect to this topology. The corresponding topological space (X, τ) is a Hausdorff locally convex space iff the family P is sufficient, that is, $\forall x_0 \in X \setminus \{\emptyset\}, \exists p_\alpha \in P$ with $p_\alpha(x_0) \neq 0$. In this context, a convex cone $K \subset X$ is an *Isac’s cone* iff $\forall p_\alpha \in P, \exists f_\alpha \in X^* : p_\alpha(x) \leq f_\alpha(x), \forall x \in K$. This is an equivalent definition of Isac’s cones in separated locally convex spaces.

The best special, refined and non-trivial Isac's cones class associated to normal cones in Hausdorff locally convex spaces was introduced and studied in (Isac, G., Bahya, A., O., 2002) as the full nuclear cones family defined as follows: if $(X, Spec(X))$ is an arbitrary locally convex space $B \subset Spec(X)$ is a Hausdorff base of $Spec(X)$ and $K \subset X$ is a normal cone, then for any mapping $\varphi : B \rightarrow K^* \setminus \{0\}$ one says that the set

$K_\varphi = \{x \in X : p(x) \leq \varphi(p)(x), \forall p \in B\}$ is the full nuclear cone associated to K whenever $K_\varphi \neq \{\theta\}$. In fact, K is supernormal iff there exists $\varphi : P \rightarrow K^* \setminus \{0\}$ such that $K \subseteq K_\varphi$ (Proposition 3.1 given by Isac, G. and Postolică, V., 2005). In the above equivalent context, if $\varphi : P \rightarrow K^*$ is a function, then the convex cone

$K_\varphi = \{x \in X : p_\alpha(x) \leq \varphi(p_\alpha)(x), \forall p_\alpha \in P\}$ is the full nuclear cone associated to K , P and φ . Taking into account that in a real normed linear space $(E, \| \cdot \|)$ a non-empty set $T \subset E$ is called a Bishop-Phelps cone if there exists y^* in the usual dual space E^* of E and $\alpha \in (0, 1]$ such that $T = \{y \in E : \alpha \|y\| \leq y^*(y)\}$ and the applications of such as these cones in Nonlinear Analysis, including the optimization programs with vector-valued objective mappings, one concludes that, for Hausdorff locally convex spaces, the full nuclear cones are natural generalizations of Bishop-Phelps cones in normed vector spaces.

The beginning and the considerations in Section 4 of (Isac, G., Bahya, A. O., 2002) suggested us to consider for each function $\varphi : P \rightarrow K^* \setminus \{0\}$ the full nuclear cone $K_\varphi = \{x \in X : p_\alpha(x) \leq \varphi(p_\alpha)(x), \forall p_\alpha \in P\}$ in order to give the next generalization of Theorem 7 in (Isac, G., Bahya, A. O., 2002), as an extension of the main result indicated by (Isac, G., Postolică, V., 2005), an important link between the efficiency, the full nuclear cones, strong optimization and a significant result concerning the multiple scalarization for this general concept of the efficiency.

Theorem 3.1. *If K is any Isac's cone in every Hausdorff locally convex space $(X, P = \{p_\alpha : \alpha \in A\})$, then*

$$(i) MIN(A, K, K_1) = \bigcup_{\substack{a \in A \\ \varphi \in P \rightarrow K^* \setminus \{0\}}} S MIN(A \cap (a - K - K_1), K_\varphi),$$

$$(ii) MAX(A, K, K_1) = \bigcup_{\substack{a \in A \\ \varphi \in P \rightarrow K^* \setminus \{0\}}} S MAX(A \cap (a + K + K_1), K_{-\varphi}),$$

$$(iii) eff(A, K, K_1) = \bigcup_{\substack{a \in A \\ \varphi \in P \rightarrow K^* \setminus \{0\}}} [S MIN(A \cap (a - K - K_1), K_\varphi) \cup S MAX(A \cap (a + K + K_1), K_{-\varphi})]$$

for any non-empty subset K_1 of K .

Proof. It follows, mutatis mutandis, the same line as for Theorem 4.3 given by Isac, G. and Postoliciă, V. in 2005. ■

Remark 3.2. To obtain (ii) from (i) it is not about of a simple formality by replacing K with $(-K)$, since $(-K)_\varphi = -K_{-\varphi}$. If $0 \notin K_1$, then $a_0 \in \text{eff}(A, K, K_1)$ implies that

$A \cap (a_0 - K - K_1) = \emptyset$. Therefore, it is not possible to have $a_0 \in S(\emptyset, K_\varphi)$. In the case of $0 \in K_1$, then $\text{eff}(A, K, K_1) = \text{eff}(A, K)$ and $a_0 \in \text{eff}(A, K)$ iff $A \cap (a_0 - K) = \{a_0\}$, so in the right member of the first proved inclusion it can be selected any convex cone, not necessary K_φ . The hypothesis $K \subseteq K_\varphi$ imposed upon the convex cone K and automatically satisfied since K is an Isac's cone was used only to prove the next inclusion, with its natural projection for $MAX(A, K, K_1)$:

$$MIN(A, K, K_1) \subseteq \bigcup_{\substack{a \in A \\ \varphi : P \rightarrow K^* \setminus \{0\}}} SMIN(A \cap (a - K - K_1), K_\varphi).$$

Corollary 3.1. For every non-empty subset A of any Hausdorff locally convex space ordered by an arbitrary Isac's cone K with its dual cone K^* we have

$$(i) MIN(A, K) = \bigcup_{\substack{a \in A \\ \varphi : P \rightarrow K^* \setminus \{0\}}} SMIN(A \cap (a - K), K_\varphi);$$

$$(ii) MAX(A, K) = \bigcup_{\substack{a \in A \\ \varphi : P \rightarrow K^* \setminus \{0\}}} SMAX(A \cap (a + K), K_{-\varphi});$$

$$(iii) \text{eff}(A, K) = \bigcup_{a \in A, \varphi \in P \rightarrow K^* \setminus \{0\}} [SMIN(A \cap (a - K), K_\varphi) \cup SMAX(A \cap (a + K), K_{-\varphi})].$$

Remark 3.3. The hypothesis of the above theorem involves K to be pointed. Consequently, $0 \in K_1$ iff $0 \in K + K_1$. If $a_0 \in S(A \cap (a - K - K_1), K_\varphi)$ for some $\varphi : P \rightarrow K^*$ and $a \in A$ with $a_0 = a - k - k_1$, $k \in K$, $k_1 \in K_1$, then $K \cap (a_0 - A) = \{0\}$ because $A \cap (a - K - K_1) \subseteq a_0 + K_\varphi$ in any such a case as this. Indeed, let $x \in K \cap (a_0 - A)$ be an arbitrary element. Then, $a_0 - x \in A$ and $a_0 - x = a - k - k_1 - x \in a - K - K_1$. Therefore, $a_0 - x \in a_0 + K_\varphi$, that is,

$-x \in K_\varphi$. For every $p_\alpha \in P$ we have $p_\alpha(-x) \leq \varphi(p_\alpha)(-x) = -\varphi(p_\alpha)(x) \leq 0$. Since p_α was arbitrary chosen in P and X is a Hausdorff locally convex space, it follows that $x=0$.

Remark 3.4. Clearly, the announced theorem represents a significant result concerning the possibilities of multiple scalarization for the study of the efficiency programs in Hausdorff separated locally convex spaces, as we can see also in the final comments of (Isac, G., Bahya, A.O., 2002) for the particular cases of Hausdorff locally convex spaces ordered by closed, pointed and normal cones.

Remark 3.5. As an open problem, it is interesting to replace K_1 with any non-empty subset of an ordered linear space X , under proper hypotheses. In the next considerations we offer significant examples and adequate remarks on the Isac's cones with the mention that for the existence of the efficient points and important properties of the efficient points sets in separated locally convex spaces ordered by (weak) such as these convex cones, through the agency of the (weak) completeness instead of compactness the reader is referred to (Isac, G., 1981, 1983, 1985, 1994, 1998; Isac, G., Postolică, V., 1993; Postolică, V., 1993, 1994, 1995, 1997, 1999, 2001, 2002, 2009; Truong, X. D. H., 1994 and the references therein).

Theorem 3.2. (Bahya, A. O., 1989). *A convex and normal cone K in a Hausdorff locally convex space is supernormal if and only if every net of K weakly convergent to zero converges to zero in the locally convex topology.*

Let us consider some pertinent examples.

1. Any pointed, convex cone, in every Euclidean space R^k ($k \in \mathbb{N}^*$) is an Isac's cone.
2. In every locally convex space any well-based convex cone is an Isac's cone.
3. A convex cone is an Isac's cone in a normed linear space if and only if it is well-based.
4. Let $n \in \mathbb{N}^*$ be arbitrary fixed and let Y be the space of all real symmetric (n, n) matrices ordered by the pointed, convex cone $C = \{A \in Y : x^T A x \geq 0, \forall x \in R^n\}$. Then, Y is a real Hilbert space with respect to the scalar product defined by $\langle A, B \rangle = \text{trace}(A \cdot B)$ for all $A, B \in Y$ and C is well-based by $B = \{A \in C : \langle A, I \rangle = 1\}$ where I denotes the identity matrix.
5. Every pointed, locally or weakly locally compact convex cone in any Hausdorff locally convex space is an Isac's cone.
6. A convex cone is an Isac's cone in a nuclear space (Pietch, A., 1972) if and only if it is a normal cone.
7. In any Hausdorff locally convex space a convex cone is an weakly Isac's cone if and only if it is weakly normal.
8. In $L^p([a, b])$, ($p \geq 1$), the convex cone $K_p = \{x \in L^p([a, b]) : x(t) \geq 0 \text{ almost everywhere}\}$ is an Isac's cone if and only if $p=1$, being well based in this case by the set $B = \{x \in K_1 : \int_a^b x(t) dt = 1\}$. Indeed, if $p > 1$, then the sequence (x_n) defined by

$$x_n(t) = \begin{cases} n^{-1/p}, & a \leq t \leq a + (b - a)/2n \\ 0, & a + (b - a)/2n < t \leq b \end{cases} \quad n \in \mathbf{N}$$

converges to 0 in the weak topology but not in the usual norm topology. Therefore, by virtue of Theorem 1, K_p is not an Isac's cone. Generally, for every $p > 1$, K_p has a base $B = \{x \in K_p : \int_a^b x(t) dt = 1\}$ which is unbounded and any cone generated by a closed and bounded set $B_t = \{x \in B : \int_a^b |x(t)|^p dt \leq t\}$ with $t \geq 0$ is certainly an Isac's cone.

A similar result holds for $L^p(\mathbf{R})$. Thus, if we consider a countable family (A_n) of Lebesgue measurable disjoint sets which covers \mathbf{R} such that $\mu(A_n) = 1$ for all $n \in \mathbf{N}$, where μ is the Lebesgue measure, then the sequence (y_n) given by $y_n(t) = 1$ if $t \in A_n$ and $y_n(t) = 0$ for $t \in \mathbf{R} \setminus A_n$ converges weakly to zero while it is not convergent to zero in the norm topology. Taking into account the above theorem, it follows that the usual positive cone in $L^p(\mathbf{R})$ is not an Isac's cone if $p > 1$, that is, it is not well-based in all these cases. However, these cones are normal for every $p \geq 1$. The same conclusion concerning the non-supernormality is valid for the positive orthant convex cones in the usual Orlicz spaces.

9. In l^p ($p \geq 1$) equipped with the usual norm $\|\cdot\|_p$ the positive cone $C_p = \{(x_n) \in l^p : x_n \geq 0 \text{ for all } n \in \mathbf{N}\}$ is also normal with respect to the norm topology, but it is not an Isac's cone excepting the case $p = 1$. Indeed, for every $p > 1$, the sequence (e_n) having 1 at the n th coordinate and zeros elsewhere converges to zero in the weak topology, but not in the norm topology and by virtue of Theorem 3 it follows that C_p is not an Isac's cone.

For $p = 1$, C_1 is well-based by the set $B = \{x \in C_1 : \|x\|_1 = 1\}$ and Proposition 5 (Isac, G., 1983) ensures that it is an Isac's cone. If we consider in this case the locally convex topology in l^1 defined by the seminorms $p_n((x_k)) = \sum_{k=0}^n |x_k|$ for every (x_k) in l^1 and $n \in \mathbf{N}$, which is weaker than its usual topology, then the usual positive cone remains an Isac's cone with respect to this topology (now it is normal in a nuclear space and one applies Proposition 6 of (Isac, G., 1983) but it is not well based. Taking into account the concept of H -locally convex space introduced by Precupanu, T. in 1969 and defined as any Hausdorff locally convex space with the seminorms satisfying the parallelogram law and the property that every nuclear space is also a H -locally convex space with respect to an equivalent system of seminorms (Pietch, A., 1972), the above example shows that in a H -locally convex space a proper convex cone may be an Isac's cone without to be well-based. Moreover, if we consider in l^2 the H -locally convex topology induced by the seminorms

$$\tilde{p}_n((x_k)) = \left(\sum_{i \geq n} |x_i|^2 \right)^{1/2}, \quad n \in \mathbf{N}, (x_k) \in \ell^2,$$

then the convex cone $C_2 = \{(x_k) \in l^2 : x_k \geq 0 \text{ for all } k \in N\}$ is normal in the H -locally convex space $(l^2, \{\bar{p}_n\}_{n \in N})$, but it is not a supernormal cone because the same sequence (e_k) is weakly convergent to zero while $(\bar{p}_n(e_k))$ is convergent to 1 for each $n \in N$ and one applies again Theorem 3. Another interesting example of normal cone in a H -locally convex space which is not supernormal is the usual positive cone in the space $L_2 \text{ loc } (R)$ of all functions from R to C which are square integrable over any finite interval of R , endowed with the system of seminorms $\{\bar{p}_n : n \in N\}$ defined by $\bar{p}_n(x) = \left(\int_{-n}^n |x(t)|^2 dt \right)^{1/2}$ for every x in $L_{loc}^2(R)$. In this case, the sequence (x_k) given by:

$$x_k(t) = \begin{cases} 0, & t \in (-\infty, 0) \cup (1/k, +\infty) \\ \sqrt{k}, & t \in [0, 1/k] \end{cases}$$

converges weakly to zero, but it is not convergent in the H -locally convex topology. The result follows by Theorem 3. It is clear that every weak topology is a H -locally convex topology and, in these cases, the supernormality of convex cones coincides with the normality thanks to the Corollary of Proposition 2 in (Isac, G., 1983).

10. In the space $C([a, b])$ of all continuous, real valued functions defined on every non-trivial, compact interval $[a, b]$ equipped with the usual supremum norm the convex cone $K = \{x \in C([a, b]) : x \text{ is concave, } x(a)=x(b) = 0 \text{ and } x(t) \geq 0 \text{ for all } t \in [a, b]\}$ is an Isac's cone, being well based by the set $\{x \in K : x(t_0) = 1\}$ for some arbitrary $t_0 \in [a, b]$. The hypothesis that all $x \in K$ are concave is essential for the supernormality.

11. The convex cone of all nonnegative sequences in the space of all absolutely convergent sequences is the dual of the usual positive cone in the space of all convergent sequences. Consequently, it has a weak star compact base and hence it is a weak star supernormal cone.

12. In l^∞ or in c_0 equipped with the supremum norm, the convex cone consisting of all sequences having all partial sums non-negative is not normal, hence it is not supernormal.

13. In every Hausdorff locally convex space any normal cone is an Isac's cone with respect to the weak topology.

14. In every locally convex lattice which is a (L) -space the ordering cone is supernormal (see also the Example 7 given by Isac, G. in 1994).

15. If we consider the space of all locally integrable functions on a locally compact space Y with respect to a Radon measure μ endowed with the topology induced by the family of seminorms $\{p_A\}$ where $p_A(f) = \int_A |f(x)| d\mu$ for every non-empty and compact subset A of Y and every locally integrable function f , then the convex cone $K = \{f : Y \rightarrow R / f(y) \geq 0, \forall y \in Y\}$ is supernormal. 16. If Z is any locally convex lattice ordered by an arbitrary convex cone K and Z^* is its topological dual ordered by the corresponding dual cone K^* , then the cone K is supernormal with respect to

the locally convex topology defined on Z by the neighbourhood base at the origin $\{[-f, f]^0\}_{f \in K^*}$.

17 In every regular vector space (E, K) (that is, the order dual E^* separates the points of E) with the property that $E = K - K$ the convex cone K is supernormal with respect to the topology defined in the preceding example.

18. Any semicomplete cone in a Hausdorff locally convex space is an Isac's cone (for this concept see the Example 11 of Isac, G., 1994).

Remark 3.6. Clearly, if a convex cone K is supernormal in a normed space, then K admits a strictly positive, linear and continuous functional, that is, there exists a linear, continuous functional f such that $f(k) > 0$ for all $k \in K \setminus \{0\}$. Generally, the converse is not true even in a Banach space as we can see in the following examples:

19. If one considers in the usual space l^p ($1 \leq p \leq \infty$) the convex cone $K = l^p_+ = \{x = (x_i) \in l^p : x_i \geq 0 \text{ for every } i \in \mathbb{N}\}$ of infinite vectors with non-negative components, then the functional φ defined by $\varphi(k) = \sum_{i=1}^{\infty} k_i$ for any $k = (k_i) \in l^p$ is linear, continuous and strictly positive. But, as we have seen in the above considerations (Example 9), this cone is supernormal if and only if $p = 1$.

20. Let K be the usual positive cone $L^p_+ = \{x \in L^p([a, b]) : x(t) \geq 0 \text{ almost everywhere}\}$ in $L^p([a, b])$ ($1 \leq p \leq \infty$). Then, the linear and continuous functional ψ on $L^p([a, b])$ given by $\psi(x) = \int_a^b x(t)dt$ for every $x \in L^p([a, b])$ is strictly positive on K while K is supernormal (see the above Example 8) if and only if $p = 1$.

Therefore, l^1_+ and L^1_+ are supernormal cones with empty topological interiors and for every $p \in (1, +\infty)$ it follows that l^p_+ and L^p_+ are normal cones with empty interiors which are not supernormal. Hence, these convex cones are not well based. A very simple example of supernormal cones having non-empty topological interior is R^n_+ ($n \in \mathbb{N}^*$).

Remark 3.7. In the order complete vector lattice $B([a, b])$ of all bounded, real valued functions on a compact non-singleton interval $[a, b]$ endowed with its usual norm the standard positive cone

$K = \{u \in B([a, b]) : u(t) \geq 0 \text{ for all } t \in ([a, b])\}$ is normal but it has not a base, that is, it is not an Isac's cone.

However, this cone has non-empty interior. If we consider the linear space l^l endowed with the separated locally convex topology generated by the family $\{p_n : n \in \mathbb{N}\}$ of seminorms defined by $p_n(x) = \sum_{k=0}^n |x_k|$ for every $x = (x_k) \in l^l$, then the convex cone $K = \{x = (x_k) \in l^l : x_k \geq 0 \text{ whenever } k \in \mathbb{N}\}$ is an Isac's cone but it is not well based.

4. SOME CONCLUSIONS

The natural context of supernormality (nuclearity) for convex cones is any separated locally convex space. Isac, G. introduced the concept of "nuclear cone" in 1981, published it in 1983 and he showed that in a normed space a convex cone is nu-

clear if and only if it is well based or equivalently iff it is “with plastering”, the last concept being defined by Krasnoselski, M. A. in fifties (see ,for example , Krasnoselski, M. A., 1964 and so on). Such a convex cone was initially called “nuclear cone” by Isac, G. (1981) because in every nuclear space (Pietch, A., 1972) any normal cone is a nuclear cone in Isac’s sense (Proposition 6 of Isac, G., 1983). Afterwards, since the nuclear cone introduced by Isac appears as a reinforcement of the normal cone, it was called supernormal. The class of supernormal cones in Hausdorff locally convex spaces was initially imposed by the theory and the applications of the efficient (Pareto minimum type) points (especially existence conditions based on completeness instead of compactness were decisive together with the main properties of the efficient points sets), the study of critical points for dynamical systems and conical support points and their importance was very well illustrated by important results, examples and comments in the specified references and in other connected papers. It is also very significant to mention again that the concept of supernormality introduced by Isac, G. (1981) is not a simple generalization of the corresponding notion defined in normed linear spaces by Krasnoselski, M. A. and his colleagues in the fifties. Thus, for example, Isac’s supernormality attached to the convex cones has his sense in every Hausdorff locally convex space identically with the well known Grothendieck’s nuclearity. By analogy with the fact that a normed space is nuclear in Grothendieck’s sense if and only if it is isomorphic with an usual Euclidean space, a convex cone is supernormal in a normed space if and only if it is well based, that is, it is generated by a convex bounded set which does not contain the origin in its closure. Beside Pareto type optimization, we also mention Isac’s significant contributions, through the agency of supernormal cones, to the convex cones in product linear spaces and Ekeland’s variational type principles (Isac, G., 2003; Isac, G., Tammer, Chr., 2003). Therefore, the more appropriate background for Isac’s cones is any separated locally convex space.

Definition 4. (Postolică, V., 1995). A nonempty set B of a topological vector space X ordered by a convex cone K is called K -bounded if there exists $B_0 \subseteq X$ bounded so that $B \subseteq B_0 \pm K$ and K -closed if its conical extension $B \pm \bar{K}$ is closed, where \bar{K} signifies the topological closure of K .

A synthesis concerning the existence and significant properties of the efficient points are given in the next theorem.

Theorem 4. (Postolică, V., 1995, 1997, 2008).

(i) if K is an Isac’s cone in a Hausdorff locally convex space X , $A, B \subseteq X$ are non-empty subsets positioned by $A \subseteq B \subseteq A + K$ ($A \subseteq B \subseteq A - K$) and $B \cap (A_0 - K)$ ($B \cap (A_0 + K)$) is a bounded and complete set for at least one non-empty set $A_0 \subseteq A$, then $MIN(A, K) \neq \emptyset$ ($MAX(A, K) \neq \emptyset$);

(ii) if X is a quasi-complete separated locally convex space, that is, any non-empty bounded and closed subset is complete and K is a closed Isac’s cone in X , then for any K -bounded and K -closed non-empty set A of X we have $MIN(A, K) \neq \emptyset$ ($MAX(A, K) \neq \emptyset$), $A \subseteq MIN_K(A) + K$
($A \subseteq MAX(A) - K$) (the domination property), $MIN_K(A) + K = A + K$

($MAX(A, K) - K = A - K$) and

$MIN(A, K)(MAX(A, K))$ is K -bounded and K -closed;

(iii) according to the same hypotheses from (ii),

$MIN(A, K) \neq \emptyset(MAX(A, K) \neq \emptyset)$ for any non-empty subset A with $B \cap (A_0 - K)(B \cap (A_0 + K))$ K -bounded and K -closed set whenever $A_0 \subseteq A \subseteq X$ and $A \subseteq B \subseteq A + K(A \subseteq B \subseteq A - K)$, with the corresponding implications for $eff(A, K)$.

Definition 5. (Postolică, V., 2008) A real function $f : X \rightarrow R$ is called $(K + K_1)$ -increasing ($K + K_1$ -decreasing) if $f(x_1) \geq f(x_2)$ whenever $x_1, x_2 \in X$ and $x_1 \in x_2 + K_1 + K(x_1 \in x_2 - K_1 - K)$.

Obviously, every real increasing (decreasing) function defined on any linear space ordered by an arbitrary convex cone K is $K + K_1$ -increasing ($K + K_1$ -decreasing), for each non-empty subset K_1 of K .

Finally, we present the next coincidence of the approximate efficient points sets and the Choquet boundaries, which generalizes the main result given by Bucur, placel. and Postolică, V., 1994, and can not be obtained as a consequence of the Axiomatic Potential Theory.

Theorem 5. (Postolică, V., 2008, 2009). If A is any non-void, compact subset of every Hausdorff locally convex space X and

(i) K is an arbitrary, closed, convex, pointed cone in X ;

(ii) K_1 is a non-empty subset of K such that $K + K_1$ is closed with respect to the separated topology on X .

Then, $MIN(A, K, K_1)(MAX(A, K, K_1))$ coincides with the Choquet boundary $\partial_{S_1}A(\partial_{S_2}A)$ of A with respect to the convex cone $S_1(S_2)$ of all $K + K_1$ -increasing ($K + K_1$ -decreasing) real continuous functions on A . Consequently, each of these sets, endowed with the corresponding trace topology, is a Baire space and a G_δ -subset of X whenever (A, τ_A) is metrizable. Moreover, $eff(A, K, K_1) = \partial_{S_1}A \cup \partial_{S_2}A$

Corollary 5.1.

(i) If for every $f \in C(A)$ one denotes $\bar{f}(a) = \sup\{f(a') : a' \in A \cap a + K + K_1\}$ and $\tilde{f}(a) = \sup\{f(a'') : a'' \in A \cap (a - K - K_1)\}$, then

$$eff(A, K, K_1) = \bigcup_{a \in A} \{a \in A : f(a) = \bar{f}(a) (\tilde{f}(a)), \forall f \in C(A)\};$$

(ii) $MIN(A, K, K_1)(MAX(A, K, K_1), MIN(A, K, K_1)((MAX(A, K, K_1)) \cap \{a \in A : s(a) \leq 0\}(s \in S_1)(s \in S_2))$ are compact sets with respect to the Choquet's topology and as usual compact subsets of A .

Remark 12. Generally, $MIN(A, K, K_1)(MIN(A, K, K_1))$ coincides with the corresponding Choquet boundary of A only with respect to the convex cone of all real, continuous and $K + K_1$ -increasing ($K + K_1$ -decreasing) functions on A . Thus, for example, if A is any non-empty, compact and convex subset of every Hausdorff locally convex space and $S = \{f : A \rightarrow R / f \text{ is continuous and concave}\}$, then its Choquet boundary with respect to S coincides with the set $ex(A)$ of all extreme points x in A , that is, if $y, z \in A$ and there exists $\alpha \in (0, 1)$ with $\alpha y + (1 - \alpha)z = x$, then $y = z = x$.

The hypothesis of concavity imposed on the functions is essential for the validity of this result. Generally, even in the finite dimensional cases, an extreme point for a compact convex set is not necessary an efficient point and conversely.

Remark 13. The largest class ζ of convex cones ensuring the existence for the efficient points in all non-empty compact subsets of every separated topological vector space was defined by Sterna – Karwat, A., 1986 as follows: if V is an arbitrary Hausdorff topological vector space, a convex cone C belongs to ζ iff for every closed vector subspace L of V , $C \cap L$ is a vector subspace whenever its closure $\overline{C \cap L}$ is a vector subspace. As we have already specified before Theorem 5, there exists more general conditions than compactness imposed upon a non-empty set A in a separated locally convex space ordered by a convex cone K ensuring that $eff(A, K) \neq \emptyset$. Perhaps our coincidence result suggests a natural extension of the Choquet boundary at least in these cases. Anyhow, Theorem 5 represents an important link between Vector Optimization and Potential Theory and a new way for the study of the properties of efficient points sets and the Choquet boundaries. Indeed, one of the main question in Potential Theory is to find the Choquet boundaries. This fact is relatively easy for particular cases but, in general, it is an unsolved problem. Since in a lot of cases the efficient points sets contain dense subsets which can be identified by adequate optimization methods, it is possible to determine the corresponding Choquet boundaries in all these situations. Consequently, our coincidence result has its practical consequences at first for the Axiomatic Theory of Potential and its applications. At the same time, by the above coincidence result, the Choquet boundaries offer important properties for the efficient points sets.

5. SOME OPEN PROBLEMS

The above context of research suggests immediately the following open problems.

Op. 5.1. If $eff(A, K) \neq \emptyset$ there exist a Hausdorff locally convex space Y , an Isac's cone K_0 in Y and a non-empty set $A_0 \subset Y$ with

$eff(A, K) = eff(A_0, K_0)$ (or, at least, $eff(A, K)$ to be dense in $eff(A_0, K_0)$)?

Op. 5.2. If $eff(A, K) \neq \emptyset$, there exist a separated locally convex space X_1 , a (pointed), convex cone K_1 in X_1 and a compact set $A_1 \subset X_1$ such that $eff(A, K) = eff(A_1, K_1)$ (at least $eff(A, K)$ to be dense in $eff(A_1, K_1)$ or conversely) ?

Op. 5.3. If T is a Hilbert space, K is a closed, convex, pointed cone in T and A is a non-empty, closed, convex subset of T does $eff(A, K)$ preserve the property of coincidence with the corresponding Choquet boundary as in the above theorem?

Op. 5.4. The same question for each of the following cases:

(i) T is a quasi-complete locally convex space, K is an Isac's cone, A is a K -bounded and K -closed set in T (Postolică, V., 1995).

(ii) T is a quasi-complete locally convex space, the closure \bar{K} of K has the properties given in (Truong, X., D. H., 1994) and A is a K -bounded and K -closed subset in T .

6. FUTURE RESEARCH DIRECTIONS

This research work can also be considered as a study in which the approximate solutions including those usual to vector optimization programs in ordered (topological) vector spaces are investigated. The listed properties of these solutions allow developments. Several new facts about the relationships between the approximate efficiency and the scalarized problems can be formulated. As we have specified, the family of Isac's cones represent the largest class of ordering cones in Hausdorff locally convex spaces ensuring the existence and the adequate properties of the efficient points sets involved in optimization, following different completeness types instead of compactness. Consequently, one of the main goal of the next research is to identify new applications of Isac's cones. The given strong connection by coincidence between the approximate solutions and Choquet's boundaries for non-empty compact sets can be extended to non-compact sets.

7. TARGET AUDIENCE

Operations Research, Multicriteria Decision Making, Mathematical Modeling.

References

- [1] R. P. Agarwal, R. P. O'REGAN, *A note on equilibria for abstract economies*, Math. Computer Modelling, 34(2001), 331 – 343.
- [2] J. P. Aubin, *Optima and equilibria. An introduction to nonlinear analysis*, Springer – Verlag Berlin Heidelberg, 1993.
- [3] A. O. Bahya, *Ensembles coniquement bornés et cônes nucléaires dans les espaces localement convexes séparés*, Thésés de Doctorat de 3-ème Cycle en Mathématiques. L'école Normale Supérieure - Takaddoum Rabat (Maroc), 1989.
- [4] H. P. Benon, – *The vector maximization problem: proper efficiency and stability*, SIAM J. Appl. Math. 32, 1977, 64-72.
- [5] H. P. Benson, *An improved definition of proper efficiency for vector minimization with respect to cones*, J. Math. Anal. Appl. 79, 1979, 232-241.
- [6] H. P. Benson, *Efficiency and proper efficiency for vector minimization with respect to cones*, J. Math. Anal. Appl. 93, 1983, 273-289.
- [7] N. Boboc, Gh. Bucur, *Convex cones of continuous functions on compact spaces*, Publishing House of Romanian Academy, Bucharest, 1976.(in Romanian)
- [8] J. M. Borwein, *Proper efficient points for maximization with respect to cones*, SIAM J. Control Optim., 15, 1977, 57 – 63.
- [9] J. M. Borwein, *On the existence of Pareto efficient points*, Math Oper. Res. 8, no. 1, 1983, 64 - 73.
- [10] I. Bucur, V. Postolică, *A coincidence result between sets of efficient points and Choquet boundaries in separated locally convex spaces*, Research Report at the 4th Workshop of the German Working Group on Decision Theory, Hotel Talblick, Holzgau, Germany, March 14-18, 1994. Optimization, 36, 1996, 231-234.

- [11] T. Cardinali, F. Papalini, *Fixed point theorems for multifunctions in topological vector spaces*, J. Math. Anal. Appl., 186, 1994, 769-777.
- [12] J. P. Dauer, R. J. Gallagher, *Positive proper efficient points and related cone results in vector optimization theory*, country-regionplaceSIAM J. Control Optim. 28(1), 1990, 158-172..
- [13] C. Dominiak, *Multicriteria decision aid under uncertainty*, In: Multiple Criteria Decision Making'05 (Ed. Tadeusz Trzaskalik), Publisher of The Karol Adamiecki University of Economics in Katowice, 2006, 63-81.
- [14] M. Ehrgott, *Multicriteria Optimization*, Springer Berlin-Heidelberg, 2005.
- [15] P. C. Fishburn, *Foundations of risk management science placeI. Risk or probable loss*, Management Science, 30, 1984, 396 – 406.
- [16] I. Gass Saul, M. Harris Carl, – *Encyclopedia of operations research and management science*, Centennial Edition, Kluwer Academic Publishers, Dordrecht, 2001.
- [17] A. M. Geoffrion, *Proper efficiency and the theory of vector maximization*, J. Math. Anal. Appl., 22, 1968, 618 -630.
- [18] R. Hartley, *On cone – efficiency, cone convexity and cone – compactness*, SIAM J. Appl. Math., 34 (2), 1978, 211- 222.
- [19] M. I. Hening, *Proper efficiency with respect to cones*, J. Optim. Theory. Appl., 36, 1982, 387-407.
- [20] P. Heyne, *Efficiency*, Research Report, University of Washington, 2000.
- [21] D. H. Hyers, G. Isac, T. M. Rassias, *Topics in nonlinear analysis and applications*, World Scientific Publishing Co, Pte. Ltd., 1997.
- [22] W. K. Kim, K.K. Tan, *New existence theorems of equilibria and applications*, Nonlinear Anal., 47(2001) 531 – 542.
- [23] G. Isac, *Cônes localement bornés et cônes complètement réguliers. Applications a l'Analyse Non-Linéaire Moderne*, Université de Sherbrooke, no. 17, 1980.
- [24] G. Isac, *Points critiques des systèmes dynamiques. Cônes nucléaires et optimum de Pareto*, Research Report, Royal Military College of St. Jean, Québec, Canada, 1981.
- [25] G. Isac, - *Sur l'existence de l'optimum de Pareto*, Riv. Mat. Univ. Parma, 4(9), 1983, 303 - 325.
- [26] G. Isac, *Supernormal cones and absolute summability*, Libertas Mathematica, 5, 1985, 17 - 32.
- [27] G. Isac, V. Postolică, - *The best approximation and optimization in locally convex spaces*, Verlag Peter Lang GmbH, Frankfurt am Main, Germany, 1993.
- [28] G. Isac, *Pareto optimization in infinite dimensional spaces: the Importance of nuclear cones*, J. Math. Anal. Appl., 182(2), 1994, 393 - 404.
- [29] G. Isac, *On Pareto efficiency. A general constructive existence principle*, Research Report (16 pp.), Department of Mathematics and Computer Science, Royal Military College of Canada, 1998.
- [30] G. Isac, A. O. Bahya, *Full nuclear cones associated to a normal cone. Application to Pareto efficiency*, Applied Mathematics Letters, 15, 2002, 633 - 639.
- [31] G. Isac, V. A. Bulavsky, V. V. Kalashnikov, *Complementarity, equilibrium, efficiency and economics*, Kluwer Academic Publishers, 2002.
- [32] G. Isac, – *Nuclear cones in product spaces. Pareto efficiency and Ekeland type variational principles in locally convex spaces* (Preprint, Royal Military College of Canada, 2003).
- [33] G. Isac, Chr. Tammer, *Nuclear and full nuclear cones in product spaces. Pareto efficiency and Ekeland type variational principles* (Preprint, Royal Military College of Canada and University of Hall, 2003).

- [34] G. Isac, V. Postolică, *Full nuclear cones and a relation between strong optimization and Pareto efficiency*, J. Global Optimiz., 32, 2005, 507 - 516.
- [35] J. Jablonsky, *A slack based model for measuring super-efficiency in data envelopment analysis*, In: Multiple Criteria Decision Making'05 (Ed.Tadeusz Trzaskalik), Publisher of The Karol Adamiecki University of Economics in Katowice, 2006, 101-112.
- [36] E. Kramar, *Locally convex topological vector spaces with hilbertian seminorms*, Rev. Roum. Math Pures et Appl., Tome XXVI, no. 1, 1981, 55-62
- [37] M. A. Krasnoselski, *Positive Solutions of Operator Equations*, Gröningen, Noordhoff, 1964.
- [38] P. Loridan, *ε -solutions of vector minimization problems*. J. Optim. Theory Appl., 43(2), 1984, 265 - 276.
- [39] D. T. Luc, *Theory of vector optimization*. Springer-Verlag, 1989.
- [40] M. Luptaeik , B.Bohm, *Measuring eco - efficiency an a Leontief input - output model*, In: Multiple Criteria Decision Making'05 (Ed.Tadeusz Trzaskalik), Publisher of The Karol Adamiecki University of Economics in Katowice, 2006, 121-135.
- [41] L. Nachbin, *Topology and order*, New York, Van Nostrand, 1965.
- [42] A. B. Nemeth, *Between Pareto efficiency and Pareto ε -efficiency*, Optimization,10(5), 1989, 615 - 637.
- [43] K. F. Ng, X. Y., Zheng, *Existence of efficient points in vector optimization and generalized Bishop-Phelps theorem*, JOTA, 115, 2002, 29 - 47.
- [44] S. Park, *Some coincidence theorems on acyclic multifunctions applications*, In: Tan, K.- K(ed) Fixed Point Theory and Applications, World Sci. NJ, 1992, 248 –277.
- [45] M. Patriche, *Fixed point theorems for nonconvex valued correspondences and applications in game theory*, Fixed point Theory, 14(2013), no. 2, 435 – 446.
- [46] A. L. Peressini, *Ordered topological vector spaces*, Harper 8 Row, New York, 1967.
- [47] A. Pierch, *Nuclear Locally Convex Spaces*, Springer – Verlag, 1972.
- [48] V. Postolică, *Spline functions in H-locally convex spaces*, An. Șt. Univ. “Al. I. Cuza”, Iași, 27, 1981, 333-338.
- [49] V. Postolică, *Existence conditions of efficient points for multifunctions with values in locally convex spaces*, Stud. Cerc. Mat., Tom 41, no. 4, 1989, 325 - 331.
- [50] V. Postolică, *New existence results for efficient points in locally convex spaces ordered by supernormal cones*, J. Global Optimiz., 3, 1993, 233 - 242.
- [51] V. Postolică, *An extension to sets of supernormal cones and generalized subdifferentials*, Optimization, 29, 1994, 131 - 139.
- [52] V. Postolică, *Properties of Pareto sets in locally convex spaces*, Optimization, 34, 1995, 223 - 229.
- [53] V. Postolică, *Recent conditions for Pareto efficiency in locally convex spaces*, Research Report at the Joint International Meeting Euro XV-INFORMS XXXIV, Barcelona, Spain, July 14 - 17, 1997.
- [54] V. Postolică, *Properties of efficient points sets and related topics*. Research Report at The Second International Conference on Multi – Objective Programming and Goal Programming, Torremolinos, Spain, May 16-18, 1996. Published in Advances in Multiple Objective and Goal Programming, Lecture Notes in Economics and Mathematical Systems, 455, (Eds.: Rafael Caballero, Francisco Ruiz, Ralph E. Steuer),1997, 201 - 209.
- [55] V. Postolică, *A method which generates splines in H-locally convex spaces and connections with vectorial optimization*, Positivity, 2 (4), 1998, 369 - 377.

- [56] V. Postolică, *Conditions for Pareto efficiency in locally convex spaces*, Mathematical Reports of Romanian Academy 2, 1(51), April-June, 1999, 257-269.
- [57] V. Postolică, A. Scarelli, *Some connections between best approximation, vectorial optimization and multicriterial analysis*, Nonlinear Analysis Forum, 5, 2000, 111 - 123.
- [58] V. Postolică, A. Scarelli, L. Venzi, *On the equilibrium of a multidimensional ecosystem*, Nonlinear Analysis Forum, 6(2), 2001, 321 - 335.
- [59] V. Postolică, *New existence results of solutions for vector optimization programs with multifunctions*, Optimization, vol. 50, 2001, 253 - 263.
- [60] V. Postolică, *Pareto efficiency, Choquet boundaries and operators in Hausdorff locally convex spaces*. Nonlinear Analysis Forum, 7(2), 2002, 215 – 230.
- [61] V. Postolică, *A class of generalized dynamical systems in connection with Pareto efficiency and related topics*, Research Paper submitted to the International Conference “Complex Systems, Intelligence and Modern Technological Applications”, Cherbourg, France, September 19 - 22, 2004. Published in the corresponding Electronics Proceedings Volume.
- [62] V. Postolică, *Choquet boundaries and efficiency*, Computer and Mathematics with Applications, 55, 2008, 381 - 391.
- [63] V. Postolică, *Approximate efficiency in infinite dimensional ordered vector spaces*, International Journal of Applied Management Science (IJAMS), Vol. 1, No. 3, 2009, 300 – 314.
- [64] T. Precupanu, *Espaces linéaires à semi-normes hilbertiennes*, An. Șt. Univ. “Al. I. Cuza”, Iași, 15, 1969, 83 - 93.
- [65] T. Precupanu, *Scalar minimax properties in vectorial optimization*, International Series of Numerical Mathematics, Birkhäuser Verlag Basel, 107, 1992, 299 - 306.
- [66] S. Songsak, W. Wing-Keung, D. Sompong, T.N.Hung, *Stochastic dominance and applications to finance, risk and economics*, CRC Press, Taylor and Francis Group, October 2009, 456 pp., ISBN: 978-1-4200-8266-1.
- [67] Z. Sonntag, C.Zălinescu, *Comparison of existence results for efficient points*, JOTA, 105, 2000, 161-188.
- [68] A.K. Sterna , *On the existence of cone maximal points in real topological linear spaces*, Israel Journal of Mathematics, 54(1), 1986, 33 - 41.
- [69] D. J. Stewart, *Uncertainties in MCDA*, In: Multi Criteria Decision Analysis (Ed. P. Greco). Springer - Verlag, 2004.
- [70] F. Trèves, *Locally convex spaces and linear partial differential equations*, Springer – Verlag, StateplaceNew York, 1967.
- [71] X. D. H. Truong, *A note on a class of cones ensuring the existence of efficient points in bounded complete sets*. Optimization, 31, 1994, 141-152.
- [72] X. D. H. Truong, *On the existence of efficient points in locally convex spaces*, J. Global Optimiz., 4, 1994, 265 - 278.
- [73] F. Wantao, C. Yonghong, *On the super efficiency in locally convex spaces*, Nonlinear Analysis 44 (2001), 821 – 828.
- [74] C. J. Zhang, *Generalized bi-quasi-variational inequalities and generalized quasi-variational inequalities*, Acta Analysis Functionalis Applicata, 7(2), 2005, 116 - 122 .
- [75] H. Zimmermann, *An application - oriented view of modeling uncertainty*, European Journal of Operation Research, 122, 2000, 190 - 198.