

EFFICIENCY IN INFINITE DIMENSIONAL ORDERED VECTOR SPACES

Vasile Postolică

Romanian Academy of Scientists,

"Vasile Alecsandri" University of Bacău, Faculty of Sciences, Department of Mathematics, Informatics and Educational Sciences,

Romania

vpostolica@ambra.ro

Abstract This research paper is focused on the common concepts of the efficiency and set-valued map. After a short introduction, we propose some questions regarding the notion of efficiency and we emphasize the Pareto optimality as one of the first finite dimensional illustrative examples. We present the efficiency and the multifunctions in the infinite dimensional ordered vector spaces following also our recent results concerning the most general concept of approximate efficiency, as a natural and the largest generalization of the efficiency, with implications and applications in vector optimization and the new links between the approximate efficiency, the strong optimization - by the full nuclear cones - and Choquet's boundaries by an important coincidence result. In this way, the efficiency is strong related to the multifunctions and Potential theory through the agency of optimization and conversely. Significant examples of Isac's cones, pertinent considerations on spline functions in H-locally convex spaces and several important references conclude this study.

Work presented as invited lecture at CAIM 2015, September 17-20, 2015, University "Ștefan cel Mare" of Suceava, Romania.

Keywords: efficiency, multifunction, Pareto optimality, optimization, approximate efficiency, Isac's (nuclear or supernormal) cone, Choquet boundary, spline function, H - locally convex space.

2010 MSC: 91C02, 90C48.

1. INTRODUCTION

Throughout in our *Life, the Efficiency* was, it is and it will remain *essential* for *the Existence* and not only. Its mathematical models were unanimously accepted in various *knowledge fields*. Also, we propose it as a *New Frontier in Mathematical Physics, in the World context of priorities concerning the Alternative Energies, the Climate Exchange and the Education*. To it we dedicate our modest scientific contribution. The content of this research work is organized as follows: *Section 2* is dedicated to some useful matters on the efficiency. We briefly present in *Section 3 the Pareto Optimality* as one of the main starting points for the mathematical modelling of the *Economical Efficiency*. In *Section 4*, conceived as a concise Survey, we present *the most general notion of Approximate Efficiency* and its particular case of

the usual *Efficiency*, with the immediate connections to *Multifunctions*, in the *Infinite Dimensional Ordered Vector Spaces*, completed by the coincidence between *Choquet's Boundaries and the Approximate Efficient Points Sets in Ordered Hausdorff Locally Convex Spaces*, this conclusion being based on the first result established by us concerning such a property as this for Pareto type efficient points sets and the corresponding Choquet boundaries. Our results represent strong relationships between important Great Fields of Mathematics and the Human Knowledge: *Vector and Strong Optimization, Set-valued Maps, the Axiomatic Theory of Potential together with its Applications and the Human Efficiency*. Since, until now, the largest class of convex cones ensuring the existence and important properties of the sets containing the efficient points following the completeness instead of compactness is represented by the Isac's (nuclear or supernormal) cones, substantial examples of such as these cones are given in Section 5, completed by the last section devoted to the spline functions in H-locally convex spaces as genuine best approximation elements and solutions for strong and vectorial optimization proper problems. Finally, we indicate the selected bibliography which refers only to the papers which were used. Sections 2 - 4 and a part of Section 5 are presented following [72].

2. SOME SELECTED QUESTIONS ON THE EFFICIENCY

First of all, we present a short survey on the efficiency. The current language defines the efficiency as “*the ability to produce the desired effect in dealing with any problem*”. In the actual world characterized by: *globalization which has generated efficiency gains, liberalisation, individualization, informatization, informalization* ([76] and so on), the efficiency is perceived as follows: “working well, quickly and without waste”. But, our life has its *Divine efficiency* and the projection in the reality deals with a lot of kind of descriptions: *information efficiency, energy efficiency, eco - efficiency* (see, as recent references, [41], [48] and so on), home energy efficiency, water efficiency market enhancement programs, sports and efficiency, economic efficiency (agricultural and industrial efficiency, human efficiency in business, efficiency in financial and betting markets, efficiency in capital asset pricing models, etc.), efficiency in Mathematics (efficient algorithms in computational complexity, efficiency frontier in data envelopment analysis, statistics and so on, efficiency in multi – objective, stochastic and goal programming, efficiency in mathematical economics, etc.), efficiency in medicine, technical efficiency, etc., all of them being based on the fundamental question: “*whose valuations do we use, and how shall they be weighted?*”[29]. Thus, the economic efficiency is characterized by the “*optimal*” relationships between the value of the ends (the physical outputs) and the value of the means (the physical inputs), both of them “measured” by the money. Hence, the monetary evaluations are essential for the economic efficiency which usually is associated with the sustainability defined as “*development that meets the needs of the present without*

compromising the ability of future generations to meet their own needs” to achieve a good society ([11], p.43). However, the goals of efficiency and sustainability might not be enough to ensure a positive social development because the economic efficiency does not include any component of it (see, for example, the distribution of goods) ([52], p.13) and the sustainability was not very clear defined [75]. In terms of the technical efficiency, a machine is considered to be useful or “*more efficient than another*” when it generates “*more work output per unit of energy input*”. But, it is not possible to speak about complete efficiency in any process since we have not developed yet workable procedures concerning the corresponding evaluations (see, for instance, the urban automobile traffic which engages different people from many points of view; in general, the transport systems are highly complex with different goals often conflict). In fact, in all the processes, we try to approximate the real efficiency by some kinds of fuzzy or relative efficiency in order to obtain several controls on it, taking into account its complexity. For example, one of the main purposes of the data envelopment analysis is to “*to measure*” the relative efficiency and the comparison of decision making units by the estimation of “*the distance*” between the evaluated real units and the virtual units (see, for instance, [40]). Another strong argument and a significant example in this direction is represented by *the permanent measure and the continuous supervision of the interest rate risk to obtain the optimal efficiency balance in the management of banks’ assets and liabilities* (see, for example, the pertinent asset liability management model together with the simulation analysis given in [42]). Generally speaking, from the decisional point of view, the main steps to obtain the efficiency are the following:

2.1. *Wording of the problems by the managerial staff in an adequate language for mathematicians and computer scientists because the “bilateral” dialogue is absolutely necessary to solve the programs through the agency of such as these cooperations.*

2.2. *The elaboration of the appropriate mathematical models and the corresponding numerical processings.*

2.3. *The selection, by a serious study, of the best multicriteria decisions.*

2.4. *The application of them.*

2.5. *The evaluation of the efficiency.*

We must remark that the multicriteria decision aid ([20], [76]) and the decision making under uncertainty viewed as an important area of decision making and recognized as “*the fact that in certain situations a person does not have the information which quantitatively and qualitatively is appropriate to describe, prescribe or predict deterministically and numerically a system, its behaviour or other characteristics*”, that is, “*it relates to a state of the human mind*” characterized by the “*lack of complete knowledge about something*” [86] represents the real risk for the efficiency which, in our opinion, generates it. Following [86], we remember that the term of “*risk*” was initially applied for “*the situations in which the probabilities of outcomes were objectively known*”. In accordance with [23] and [77] now this concept “*means*

a possibility of something bad happening” and the uncertainty “is applied to the problems in which real alternatives with several possible outcomes exist”. Another important matter is the efficacy defined as “the quality of being efficacious” in the sense of “producing the desired effects or results”. In our opinion, this means to be efficient step by step, that is, a discrete efficiency with appropriate links between the stages.

3. PARETO OPTIMALITY: AN ILLUSTRATIVE EXAMPLE OF EFFICIENCY

Not even for the market economies there exists no an universal mathematical model. Pareto efficiency or Pareto optimality [54] is a central theory in economics with broad applications in game theory, engineering and the social sciences. It represents the actual finite dimensional part of the multiobjective programming in vector optimization. Thus, whenever a feasible deviation from a genuine solution S of an arbitrary multiobjective programme generates the improvement of at least one of the objectives while some other objectives degrade, any such a solution S as this is called efficient or nondominated. A system in economics and in politics is called Pareto efficient whenever “no individual can be made better off without another being made worse off” that is, a social state is economically efficient, or Pareto optimal, provided that “no person in society can become better off without anyone else becoming worse off”. This characteristic of Pareto type efficiency has been pointed out in [1] - [3], [19], [26], [54], [80] and the others. In terms of the alternative allocations this means that given a set of alternative allocations and a set of individuals, any movement from one alternative allocation to another that can make at least one individual better off, without making any other individual worse off is called a Pareto improvement or a Pareto optimization. An allocation of resources is named *Pareto efficient* or *Pareto optimal* whenever *no further Pareto improvements can be made*. If the allocation is *strictly preferred* by one person and *no other allocation would be as good for everyone*, then it is called *strongly Pareto optimal*. A *weakly Pareto optimal allocation* is one where *any feasible reallocation would be strictly preferred by all agents* [86]. Consequently, *Pareto type efficiency is an important approximate criterion for evaluating the economic systems and the political policies, with minimum assumptions on the interpersonal comparability*. We said “*approximate criterion*” because it asks “*the ideal*” which may not reflect, for example, the workings of real economics, thanks to the following restrictive assumptions necessary for the existence of Pareto efficient outcomes: the markets exist for all possible goods, being perfectly competitive and the transaction costs are negligible. In the political policies, not every Pareto efficient outcome is regarded as desirable (see, for instance, the strategies based on the unilateral benefits). For these reasons, Pareto optimality was accepted with some or much uncertainty and controversy, but, by the Arrow’s renowned impossibility theorem given in 1951 according to which “*no social preference ordering based on*

individual orderings only could satisfy a small set of very reasonable conditions”- the Pareto criterion being one of them, it remains ”plausible and uncontroversial” [75]. Usually, the concept of efficiency replaces the notion of optimality in multiple criteria optimization because *whenever the solutions of a multiple – objectives program exist in Pareto’sense, they cannot be improved following the ordering induced by the cone.*

4. APPROXIMATE EFFICIENCY, EFFICIENCY AND MULTIFUNCTIONS IN INFINITE DIMENSIONAL ORDERED VECTOR SPACES WITH RECENT CONNECTED RESULTS

Seemingly, the concept of efficiency is equivalent to the optimality, as we can see from the next abstract construction. In reality, the optimality represents a particular case of the efficiency, that is, “the best approximation” of all the efficient points. Let X be a real or complex ordered vector space, let \mathcal{K} be the class of all convex cones defined on X and let A be an arbitrary, non-empty subset of X . Following the next considerations, we consider that the set of all efficient points of A with respect to an arbitrary $K \in \mathcal{K}$ is in the following relation with the “vectorial” minimization or maximization:

$$Eff(A) \supseteq \bigcup_{K \in \mathcal{K}} MIN_K(A) \bigcup \bigcup_{K \in \mathcal{K}} MAX_K(A).$$

Clearly, any vector optimization program (which has its origin in the usual ordered Euclidean spaces programs thanks to Pareto optimality of vector-valued real functions) includes the corresponding strong optimization program and allows to be described as one of the next optimization problems. Moreover, it is possible to replace the convex cone K by any other convenient, non-empty subset of X :

$$(P_{T,K}) : MIN_K f(T) \text{ or } (P_{T,K}) : MAX_K f(T)$$

where $K \in \mathcal{K}$, T is a given non-empty set and $f : T \rightarrow X$ is an appropriate application. If one denotes by $S_f(T, K)$ the corresponding set of solutions, then the announced equivalence seems to be justified by the following relation:

$$\bigcup_{\emptyset \neq A \subseteq X} Eff(A) = \bigcup_{\substack{T \neq \emptyset \\ K \in \mathcal{K}, f : T \rightarrow X}} S_f(T, K)$$

but, in reality, only the next inclusion is valid :

$$\bigcup_{\emptyset \neq A \subseteq X} Eff(A) \supseteq \bigcup_{\substack{T \neq \emptyset \\ K \in \mathcal{K}, f : T \rightarrow X}} S_f(T, K)$$

thanks to the refinement of the notion of efficiency which can not be totally described for the moment by mathematical means, that is, the final agreement between “the dictionary’s concept of efficiency” and “the corresponding notion in mathematics” was not signed yet.

Now, let X be a non-empty set, let E be a vector space ordered by a convex, pointed cone K and let $f : X \rightarrow E$ be a function. We consider the next Vector Optimization Problem:

$$(P) \begin{cases} \min f(x) \\ x \in X \end{cases},$$

To solve it means to identify all the efficient points $x_0 \in X$ in the sense that $f(X) \cap [f(x_0) - K] = \{f(x_0)\}$. In all these cases, x_0 is an *efficient solution* and $f(x_0)$ is called a *nondominated point* of the program (P) for Multicriteria Optimization Programs in Finite Dimensional Vector Spaces. If one replaces K by $K \setminus \{0\}$ or $\text{int}(K)$, then x_0 is called *strictly (weakly) efficient* for X , respectively, and $f(x_0)$ is named *strictly (weakly) nondominated*. Consequently, whenever E is an usual ordered Euclidean space, the above concepts for x_0 means *Pareto optimal*, *strictly Pareto optimal* and *weakly Pareto optimal solution*, respectively, with the corresponding nondominated concepts for f (see, for a recent instance, [21]). So, concerning the practical vector - optimization problems, it is important to know when the set of all efficient points is non-empty, to establish its main properties (existence, domination, connectedness, compactness, density in varied topologies, etc.) and to extend the concepts together with the results to multicriteria optimization in infinite dimensional ordered vector spaces. The proper efficiency introduced in [45] and developed in [5] – [7], [9], [10], [18], [25], [27], [28], [35], [38], [50], etc. appears as a refined case of the efficiency, that is, the set of all properly efficient solutions and the set of all positive proper efficient solutions of problem (P) are subsets of the set containing all the efficient points and the study of them has been proposed in order to eliminate some “undesirable” efficient solutions. Moreover, in the recent research work [22], the author propose a slight modification to the well known *Geoffrion proper efficiency* in real *vector optimization* with the main aim to generalize some substantial results in the case of *infinitely many criteria*, multicriteria approaches for *decision-making* under uncertainty and *stochastic optimization*. We recall that $x_o \in X$ is a *properly efficient solution* of (P) if it is an *efficient point* and $\text{cl}[\text{cone}(f(X) + K - \{f(x_o)\})] \cap K = \{0\}$; x_o is a *positive proper efficient solution* of (P) if there exists a linear continuous functional φ on E such that $\varphi(k) > 0$ for every $k \in K$ and $\varphi[f(x_o)] \leq \varphi[f(x)]$ for all $x \in X$. This section deals with a new generalization of the efficiency, named by us the *approximate efficiency*, in ordered Hausdorff locally convex spaces. All the elements concerning the ordered topological vector spaces used here are in accordance with [55].

Let E be a vector space ordered by a convex cone K, K_1 a non-void subset of K and A a non-empty subset of E . The following definition introduces a new concept of approximate efficiency which generalizes the well known notion of Pareto efficiency.

Definition 1.[71]. We say that $a_0 \in A$ is a K_1 -Pareto (minimal) efficient point of A , in notation, $a_0 \in \text{eff}(A, K, K_1)$ (or $a_0 \in \text{MIN}_{K+K_1}(A)$) if it satisfies one of the following equivalent conditions:

- (i) $A \cap (a_0 - K - K_1) \subseteq a_0 + K + K_1$;
- (ii) $(K + K_1) \cap (a_0 - A) \subseteq -K - K_1$;

In a similar manner one defines the Pareto (maximal) efficient points by replacing $K + K_1$ with $-(K + K_1)$. Clearly,

$$A \cap (a_0 - K) \subseteq a_0 + K_1 \Rightarrow A \cap (a_0 - K - K_1) \subseteq a_0 + K + K_1 \Rightarrow A \cap (a_0 - K_1) \subseteq a_0 + K,$$

which suggests other concepts for the approximate efficiency in ordered linear spaces.

Remark 1. The notion of *approximate efficiency* in the sense of the above definition is *the most general notion of approximate efficiency* introduced until now. We also remark that $a_0 \in \text{eff}(A, K, K_1)$ iff it is a *fixed point* for the multifunction $F : A \rightarrow A$ defined by

$$F(t) = \{a \in A : A \cap (a - K - K_1) \subseteq t + K + K_1\}.$$

Consequently, for the existence of the Pareto type efficient points we can apply appropriate fixed points theorems for multifunctions (see, for instance, [13], [53], [85] and any other proper scientific papers) and we need the next usual continuity properties of multifunctions. Namely, if X and Y are two topological spaces and $f : X \rightarrow 2^Y$ is a set-valued map, then: f is *upper semicontinuous (usc)* if for any $x \in X$ and any open set $D \supseteq f(x)$ we have $D \supseteq f(x_1)$ for all x_1 in some neighbourhood $V(x)$ of x . The multifunction f is *lower semicontinuous (lsc)* if for any $x \in X$ and every open set D with $f(x) \cap D \neq \emptyset$ it follows that $f(x_1) \cap D \neq \emptyset$ for all x_1 from some neighbourhood $V(x)$ of x . The map f is *continuous* iff it is both usc and lsc. Whenever *the graph* of f defined by $\text{gr}(f) = \{(x, y) \in X \times Y : y \in f(x)\}$ is a *closed (open)* set of $X \times Y$ one says that f has a closed (open) graph. Any multifunction having a closed graph is also called *closed*. If $f(x)$ is a closed (compact) subset of Y for any $x \in X$, then f is a *closed-valued (compact-valued)* map. The set-valued map f is called *compact* if $\text{im}(f) = f(X) = \bigcup_{x \in X} f(x)$ is contained in a compact subset of Y . A topological space is *acyclic* if all of its reduced Eech homology groups over rationals vanish (see for example, the contractible spaces; in particular, the convex sets and the star-shaped sets are acyclic). The multifunction f is *acyclic* if it is usc and $f(x)$ is non-empty, compact and acyclic for all $x \in X$. The next theorem is useful to establish the existence of the (approximate) efficient points taking into account the above mentioned connection with the fixed points of multifunctions.

Theorem 1. [53]. *If A is any non-empty convex subset of an arbitrary Hausdorff separated locally convex space and $F : A \rightarrow 2^A$ a compact acyclic multifunction, then F has a fixed point, that is, there exists $a_0 \in A$ such that $a_0 \in F(a_0)$.*

Remark 2. In [50] it was proved that whenever $K_1 \subset K \setminus \{0\}$, the existence of this new type of efficient points for bounded from below sets characterizes the semi-Archimedean ordered vector spaces and the regular ordered locally convex spaces.

Remark 3. When K is pointed, that is, $K \cap (-K) = \{0\}$, then $a_0 \in \text{eff}(A, K, K_1)$ means that $A \cap (a_0 - K - K_1) = \emptyset$ or, equivalently, $(K + K_1) \cap (a_0 - A) = \emptyset$ for $0 \notin K_1$ and $A \cap (a_0 - K - K_1) = \{a_0\}$, respectively, if $0 \in K_1$. Whenever $K_1 = \{0\}$, from Definition 1 one obtains the usual concept of *efficient (Pareto minimal, optimal or admissible) point in the usual Euclidean ordered spaces* : $a_0 \in \text{eff}(A, K)$

(or $a_0 \in \text{MIN}_K(A)$) if it fulfils (i), (ii) or any of the next equivalent properties:

$$(iii) (A+K) \cap (a_0 - K) \subseteq a_0 + K ;$$

$$(iv) K \cap (a_0 - A - K) \subseteq -K$$

These relations show that a_0 is a *fixed point* for each of the following *multifunctions*:

$$F_1 : A \rightarrow A, F_1(t) = \left\{ \alpha \in A : A \cap (\alpha - K) \subseteq t + K \right\},$$

$$F_2 : A \rightarrow A, F_2(t) = \left\{ \alpha \in A : A \cap (t - K) \subseteq \alpha + K \right\},$$

$$F_3 : A \rightarrow A, F_3(t) = \left\{ \alpha \in A : (A + K) \cap (\alpha - K) \subseteq t + K \right\},$$

$$F_4 : A \rightarrow A, F_4(t) = \left\{ \alpha \in A : (A + K) \cap (t - K) \subseteq \alpha + K \right\},$$

that is, $a_0 \in F_i(a_0)$ for every $i = \overline{1,4}$. If, in addition, K is pointed, then $a_0 \in A$ is an efficient point of A with respect to K if and only if one of the following equivalent relations holds:

$$(v) A \cap (a_0 - K) = \{a_0\};$$

$$(vi) A \cap (a_0 - K \setminus \{0\}) = \emptyset;$$

$$(vii) K \cap (a_0 - A) = \{0\};$$

$$(viii) (K \setminus \{0\}) \cap (a_0 - A) = \emptyset;$$

$$(ix) (A + K) \cap (a_0 - K \setminus \{0\}) = \emptyset.$$

and we notice that $\text{eff}(A, K) = \bigcap_{\{0\} \neq K_2 \subseteq K} \text{eff}(A, K, K_2)$. Moreover, $a_0 \in \text{eff}(A, K)$ iff it is a *critical (equilibrium) point* ([31], [32], [67], [70]) for the *generalized dynamical system* $\Gamma : A \rightarrow 2^A$ defined by $\Gamma(a) = A \cap (a - K)$, $a \in A$. Thus, $\text{eff}(A, K)$ describes the moments of equilibrium for Γ and *the ideal equilibria* are contained in this set. Taking $K_1 = \{\varepsilon\} (\varepsilon \in K \setminus \{0\})$, it follows that $a_0 \in \text{eff}(A, K, K_1)$ iff $A \cap (a_0 - \varepsilon - K) = \emptyset$. In all these cases, the set $\text{eff}(A, K, K_1)$ is denoted by $\varepsilon\text{-eff}(A, K)$ and it is obvious that $\text{eff}(A, K) = \bigcap_{\varepsilon \in K \setminus \{0\}} [\varepsilon\text{-eff}(A, K)]$. Concerning existence results on the efficient points and significant properties for the efficient points sets we refer the reader to [12], [31] – [37], [46], [47], [50], [51], [59] – [70], [79], [82], [83] for a survey. The following theorem offers the first immediate connection between the strong optimization and this kind of approximate efficiency, in the environment of the ordered vector spaces.

Theorem 2. [39]. *If we denote by $S(A, K, K_1) = \{a_1 \in A : A \subseteq a_1 + K + K_1\}$ and $S(A, K, K_1) \neq \emptyset$, then $S(A, K, K_1) = \text{eff}(A, K, K_1)$.*

Remark 4. We shall denote by $S(A, K, \{0\})$ the set $S(A, K, \{0\})$. If $S(A, K, K_1) \neq \emptyset$, then $K + K_1 = K$, hence $\text{eff}(A, K, K_1) = \text{eff}(A, K)$. Indeed, let $a \in S(A, K, K_1)$. Then, $a \in a + K + K_1$ which implies that $0 \in K + K_1$. Therefore, $K \subseteq K_1 + K + K = K_1 + K \subseteq K$.

The above theorem shows that, for any non-empty subset of an arbitrary vector space, the set of all strong minimal elements with respect to any convex cone through the agency of every non-noid subset of it coincides with the corresponding set of the efficient points, whenever there exists at least a strong minimal element. Obviously, the result remains valid for the strong maximal elements and the corresponding efficient points, respectively.

Using this conclusion and the abstract construction given in [57] and [64] for the splines in the H-locally convex spaces introduced in [73] as separated locally convex spaces $(X, P = \{p_\alpha : \alpha \in I\})$ with any semi-norm $p_\alpha (\alpha \in I)$ satisfying the parallelogram law: $p_\alpha^2(x+y) + p_\alpha^2(x-y) = 2[p_\alpha^2(x) + p_\alpha^2(y)]$ whenever $x, y \in X$, linear topological spaces also studied in [43], it follows that the only best simultaneous and vectorial approximation for each element in the direct sum of any (closed) linear subspace and its orthogonal, with respect to any linear (continuous) operator between two arbitrary H-locally convex spaces, is its spline function. We also note that it is possible to have $S(A, K, K_1) = \emptyset$ and $\text{eff}(A, K, K_1) = A$. Thus, for example, if one considers $X = R^n (n \in N, n \geq 2)$ endowed with the separated H - locally convex topology generated by the semi-norms $p_i : X \rightarrow R_+, p_i(x) = |x_i|, \forall x = (x_i) \in X, i = 1, n, K = R_+^n, K_1 = \{(0, \dots, 0)\}$ and for each real number c we define $A_c = \{(x_i) \in X : \sum_{i=1}^n x_i = c\}$, then it is clear that $S(A_c, K, K_1)$ is empty and $\text{eff}(A_c, K, K_1) = A_c$.

At the same time, in the usual real linear space of all sequences, ordered by the convex cone $K = \{(x_n) : n \in N^*, n \geq 2, x_n \geq 0, \forall n \geq 2\}$, for $A = \{(x_{n\alpha}) : n \in N^*, n \geq 2, \alpha > 0\}$ with $x_{n\alpha} = (n-1)^{-\alpha} - n^{-\alpha}, n \in N^*, n \geq 2, \alpha > 0$ and $K_1 = \{(0, 0, \dots)\}$ we have

$$\text{Eff}(A, K, K_1) = S(A, K, K_1) = \emptyset.$$

In all our further considerations we suppose that X is a Hausdorff Locally Convex Space having the topology induced by a family $P = \{p_\alpha : \alpha \in I\}$ of semi-norms, ordered by a convex cone K and its topological dual space X^* . In this framework, the next theorem contains a significant criterion for the existence of the approximate efficient points, in particular, for the usual efficient points, taking into account that the dual cone of K is defined by $K^* = \{x^* \in X^* : x^*(x) \geq 0, \forall x \in K\}$ and its attached polar cone is $K^0 = -K^*$. The cone K is called (nuclear or supernormal) [31], [32] and Isac's cone [71] if for every seminorm $p_\alpha \in P$ there exists $f_\alpha \in X^*$ such that $p_\alpha(k) \leq f_\alpha(k)$ for all $k \in K$. Moreover, if $\varphi : P \rightarrow K^*$ is a function, then the convex cone $K_\varphi = \{x \in X : p_\alpha(x) \leq \varphi(p_\alpha)(x), \forall p_\alpha \in P\}$ is the full nuclear cone associated

to K , P and φ [37]. A characterization of the supernormality by the full nuclearity is given in the next Remark 7.

Theorem 3. [39]. *If A is any non-empty subset of X and K_1 is every non-void subset of K , then $a_0 \in \text{eff}(A, K, K_1)$ whenever for each $p_\alpha \in P$ and there exists x^* in the polar cone K^0 of K such that $p_\alpha(a_0 - a) \leq x^*(a_0 - a) + \eta$, $\forall a \in A$.*

Remark 5. The above theorem represents an immediate extension of Proposition 1.2 in [74]. Generally, the converse of this theorem is not valid at least in partially ordered separated locally convex spaces as we can see from the example considered in Remark 4. Indeed, if one assumes the contrary in the corresponding mathematical background then, taking $\eta = \frac{1}{4}$, it follows that for each $\lambda_0 \in [0, 1]$, there exists $c_1, c_2 \leq 0$ such that $|\lambda_0 - \lambda| \leq (c_1 - c_2)(\lambda_0 - \lambda) + \frac{1}{4}$, $\forall \lambda \in [0, 1]$.

Taking $\lambda_0 = \frac{1}{4}$ one obtains $|1 - 4\lambda| \leq (c_1 - c_2)(1 - 4\lambda) + 1$, $\forall \lambda \in [0, 1]$ which for $\lambda = 0$ implies that $c_2 \leq c_1$ and for $\lambda = \frac{1}{2}$ leads to $c_1 \leq c_2$, that is, $|1 - 4\lambda| \leq 1$, $\forall \lambda \in [0, 1]$, a contradiction.

Remark 6. If $a_0 \in A$ and for every $p_\alpha \in P$, $\eta \in (0, 1)$ there exists $x^* \in K^0$ such that $p_\alpha(a_0 - a) \leq x^*(a_0 - a) + \eta$, $\forall a \in A$, then $K \cap (a_0 - A) = \{0\}$ even if K is not pointed. Indeed, if $x \in K \cap (a_0 - A)$, then $a_0 - x \in A$ and for each $p_\alpha \in P$ and $\eta \in (0, 1)$ there exists $x^* \in K^0$ with $p_\alpha(x) = p_\alpha(a_0 - (a_0 - x)) \leq x^*(x) + \eta \leq \eta$. Because η is arbitrarily chosen in $(0, 1)$, we obtain $p_\alpha(x) = 0$ and since X is separated it follows that $x = 0$. If $0 \in K + K_1$, then $K + K_1 = K$ and $0 \notin K + K_1$ implies that $(K + K_1) \cap (a_0 - A) = \emptyset$. Consequently, $a_0 \in \text{eff}(A, K, K_1)$ in both cases and in this way we indicated also another proof of the theorem. The beginning and the considerations in Section 4 of [37] suggested us to consider for each function $\varphi : P \rightarrow K^* \setminus \{0\}$ the full nuclear cone

$K_\varphi = \{x \in X : p_\alpha(x) \leq \varphi(p_\alpha)(x), \forall p_\alpha \in P\}$ in order to give the next generalization of Theorem 7 indicated in [36].

Theorem 4. [39]. *If $0 \in K_1$ and there exists $\varphi : P \rightarrow K^* \setminus \{0\}$ with then*

$$\text{eff}(A, K, K_1) = \bigcup_{\substack{a \in A \\ \varphi \in P \rightarrow K^* \setminus \{0\}}} S(A \cap (a - K - K_1), K_\varphi)$$

for any non-empty subset K_1 of K .

Remark 7. If $0 \notin K_1$, then $a_0 \in \text{eff}(A, K, K_1)$ implies that $A \cap (a_0 - K - K_1) = \emptyset$. Therefore, it is not possible to have $a_0 \in S(\emptyset, K_\varphi)$. In case of $0 \in K_1$, then $\text{eff}(A, K, K_1) = \text{eff}(A, K)$ and $a_0 \in \text{eff}(A, K)$ iff $A \cap (a_0 - K) = \{a_0\}$, so in the right member of the first proved inclusion it can be selected any convex cone, not necessary K_φ . The hypothesis $K \subseteq K_\varphi$ imposed upon the convex cone K is automatically satisfied whenever K is an Isac's (nuclear or supernormal) cone ([31] – [35], [37]) and it was used only to prove the inclusion

$$\text{eff}(A, K, K_1) \subseteq \bigcup_{a \in A, \varphi : P \rightarrow K^* \setminus \{0\}} S(A \cap (a - K - K_1), K_\varphi).$$

Moreover, K is an Isac's cone if and only if there exists $\varphi : P \rightarrow K^* \setminus \{0\}$ such that $K \subseteq K_\varphi$. Indeed, Lemma 5 of [37] ensures the necessity of the above inclusion condition. Conversely, since for every seminorm $p_\alpha \in P$ there exists $\varphi(p_\alpha) \in K^* \setminus \{0\}$ and for any $x \in K \subseteq K_\varphi$ it follows that $p_\alpha(x) \leq \varphi(p_\alpha)(x)$, we conclude the nuclearity of K . When K is an arbitrary pointed convex cone, A is a non-empty subset of X and $a_0 \in \text{eff}(A, K)$, then, by virtue of (v) in Remark 3, we have $A \cap (a_0 - K) = \{a_0\}$, that is, $A \cap (a_0 - K) - a_0 = \{0\} \subset K_\varphi$. Hence, $a_0 \in S(A \cap (a_0 - K), K_\varphi)$ for every mapping $\varphi : P \rightarrow K^* \setminus \{0\}$ and the next corollary is valid.

Corollary 4.1. *For every non-empty subset A of any Hausdorff locally convex space ordered by an arbitrary, pointed convex cone K with its dual cone K^* we have*

$$\text{eff}(A, K) = \bigcup_{\substack{a \in A \\ \varphi : P \rightarrow K^* \setminus \{0\}}} S(A \cap (a - K), K_\varphi)$$

Remark 8. The hypothesis of Theorem 3 together with Lemma 3 of [37] involves K to be pointed. Consequently, $0 \in K_1$ iff $0 \in K + K_1$. If $a_0 \in S(A \cap (a - K - K_1), K_\varphi)$ for some $\varphi : P \rightarrow K^*$ and $a \in A$ with $a_0 = a - k - k_1$, $k \in K$, $k_1 \in K_1$, then $K \cap (a_0 - A) = \{0\}$ because $A \cap (a - K - K_1) \subseteq a_0 + K_\varphi$ in any such a case as this. Indeed, let $x \in K \cap (a_0 - A)$ be an arbitrary element. Then, $a_0 - x \in A$ and $a_0 - x = a - k - k_1 - x \in a - K - K_1$. Therefore, $a_0 - x \in a_0 + K_\varphi$, that is, $x \in K_\varphi$. For every $p_\alpha \in P$ we have $p_\alpha(-x) \leq \varphi(p_\alpha)(-x) = -\varphi(p_\alpha)(x) \leq 0$. Since p_α was arbitrary chosen in P and X is a Hausdorff locally convex space, it follows that $x = 0$.

Remark 9. Clearly, the announced theorem represents a significant result concerning the possibilities of scalarization for the study of some Pareto efficiency programs in separated locally convex spaces, as we can see also in the final comments of [37] for the particular cases of Hausdorff locally convex spaces ordered by closed, pointed and normal cones.

Remark 10. As an open problem, it is interesting to replace K_1 with any non-empty subset of an ordered linear space X , under proper hypotheses.

Remark 11. It is well known that the Choquet boundary represents a basic concept in the axiomatic theory of potential and its applications and the efficiency is a fundamental notion in vector optimization. The main aim of the last part in this section is to indicate the recent generalization of our coincidence result established in [12] between the set of all Pareto type minimum points of any non-empty, compact set in an ordered Hausdorff locally convex space and the Choquet boundary of the same set with respect to the convex cone of all real, increasing and continuous functions defined on the set, using our new concept of approximate efficiency. Following this line, firstly the Choquet boundary concept is revised in an original manner.

Let us consider an arbitrary Hausdorff locally space (E, τ) , where τ denotes its topology and let K be any closed, convex, pointed cone in E . The usual order relation \leq_K associated with K is defined by $x \leq_K y$ ($x, y \in E$) if there exists $k \in K$ with $y = x + k$. Clearly, this order relation on E is closed, that is, the set G_K given by

$G_K = \{(x, y) \in E \times E : x \leq_K y\}$ is a closed subset of $E \times E$ endowed with the induced product topology. If S is any convex cone satisfying the properties:

a) $\forall x \in X, \exists s \in S, s > 0$ and $s(x) < +\infty$;

b) S linearly separates $X_1 = \{x \in X : \exists s \in S \text{ with } s(x) < 0\}$, that is, for every $x, y \in X_1, x \neq y$, there exists $s, t \in S$ with real values in x and y such that $s(x)t(y) \neq s(y)t(x)$, then, on the set $M_+(X)$ of all positive Radon measures defined on X , one associates the following natural pre-order relation: if $\mu, \nu \in M_+(X)$, then $\mu \leq_S \nu$ means that $\mu(s) \leq \nu(s)$ for all $s \in S$.

Let S_1 be the convex cone of all lower semicontinuous and bounded from below real functions on X having the next property: if $x \in X$ and $\mu \leq_S \varepsilon_x$, where $\varepsilon_x(f) = f(x)$ for every real continuous function f on X denotes the Dirac measure, implies that $\mu(s) \leq s(x)$. Any non-empty subset $T \subseteq X$ will be called

S -boundary if, whenever $s \in S_1$ and its restriction on T denoted by $s|_T$ is positive, it follows that $s \geq 0$. The smallest, closed

S -boundary is usually called *the Silov boundary of X with respect to S* . A closed set $A \subseteq X$ is called S -absorbent if $x \in A$ and $\mu \leq_S \varepsilon_x$ implies that $\mu(X \setminus A) = 0$. The set $\partial_S X = \{x \in X_1 : \{x\} \text{ is } S\text{-absorbent}\}$ is named *the Choquet boundary of X with respect to S* and clearly its closure coincides with *the Silov boundary of X with respect to S* . The trace on $\partial_S X$ of the topology on X in which the closed sets coincide with X or with any of the

S -absorbent subsets of X contained in X_1 is usually called *the Choquet topology of $\partial_S X$* .

Definition 2. [71]. A real function $f : X \rightarrow R$ is called $(K + K_1)$ -increasing if $f(x_1) \geq f(x_2)$ whenever $x_1, x_2 \in X$ and $x_1 \in x_2 + K_1 + K$.

It is obvious that every real increasing function defined on any linear space ordered by an arbitrary convex cone K is $K + K_1$ -increasing, for each non-empty subset K_1 of K .

Now, we present the coincidence of the approximate efficient points sets and the Choquet boundaries, which generalizes the main results given in [11] and [61], respectively, and can not be obtained as a consequence of the Axiomatic Potential Theory.

Theorem 5. [71]. If A is any non-void, compact subset of X and

(i) K is an arbitrary, closed, convex, pointed cone in X ;

(ii) K_1 is a non-empty subset of K such that $K + K_1$ is closed with respect to the Hausdorff separated locally convex topology on X .

Then, $eff(A, K, K_1)$ coincides with the Choquet boundary of A with respect the convex cone S_1 of all $K + K_1$ -increasing real continuous functions on A . Consequently, the set $eff(A, K, K_1)$ endowed with the corresponding trace topology is a Baire space and, if (A, τ_A) is metrizable, then $eff(A, K, K_1)$ is a

G_δ -subset of X .

Corollary 5.1.

(i) $eff(A, K, K_1) = \{a \in A : f(a) = \sup \{f(a') : a' \in A \cap (a - K - K_1)\} \forall f \in C(A)\}$;

(ii) $eff(A, K, K_1)$ and $eff(A, K, K_1) \cap \{a \in A : s(a) \leq 0\}$ ($s \in S$) are compact sets with

respect to Choquet's topology;

(iii) $eff(A, K, K_1)$ is a compact subset of A .

Corollary 5.2. *Under the hypotheses of the above theorem we proved that*

$$eff(A, K, K_1) = \{a \in A : f(a) = \bar{f}(a), \forall f \in C(A)\} = \partial_{S_1} A = \{a \in A : a \in F(a)\}$$

where $F : A \rightarrow A$ is the multifunction defined by

$$F(t) = \left\{ a \in A : A \bigcap (a - K - K_1) \subseteq t + K + K_1 \right\}, \forall t \in A.$$

Remark 12.

In general, $eff(A, K, K_1)$ coincides with the Choquet boundary of A only with respect to the convex cone of all real, continuous and $K + K_1$ - increasing functions on A . Thus, for example, if A is a non-empty, compact and convex subset of X , then the Choquet boundary of A with respect to the convex cone of all real, continuous and concave functions on A coincides with the set of all extreme points for A . But, it is easy to see that, even in finite dimensional cases, an extreme point for a compact convex set is not necessary an efficient point and conversely.

Remark 13.

As we have already specified before Theorem 1, there exists more general conditions than compactness imposed upon a non-empty set A in a separated locally convex space ordered by a convex cone K ensuring that $eff(A, K) \neq \emptyset$. Perhaps our coincidence result suggests a natural extension of the Choquet boundary at least in these cases. Anyhow, Theorem 4 represents an important link between vector optimization and potential theory and a new way for the study of the properties of efficient points sets and the Choquet boundaries. Indeed, one of the main question in potential theory is to find the Choquet boundaries. This fact is relatively easy for particular cases but, in general, it is an unsolved problem. Since in a lot of cases the efficient points sets contain dense subsets which can be identified by adequate optimization methods, it is possible to determine the corresponding Choquet boundaries in all these situations. In this direction of study, an important role is attributed to the density properties of the efficient points sets with respect to varied topologies. Consequently, our coincidence result has its practical consequences at first for the axiomatic theory of potential and its applications. At the same time, by the above coincidence result, the Choquet boundaries offer important properties for the efficient points sets. In this way, the above coincidence result establishes a strong relationship between the approximate (in particular, strong) solutions for vector optimization programs in separated, ordered topological vector spaces and Choquet's boundaries of non-empty compact sets. Similar to the Choquet integral considered as an important risk measure [77], Choquet's boundary represents a very significant efficiency mathematical

model. Both of these last notions were introduced by Choquet in different moments of his life, so *there exists no any connection* between them.

5. EXAMPLES OF ISAC'S (SUPERNORMAL) CONES

In the next considerations we offer *significant examples* and *adequate remarks* on *the nuclear or supernormal cones*. ([31],[32]) named by us in [71] *Isac's cone* The existence of the efficient points and important properties of the efficient points sets are ensured in separated locally convex spaces ordered by (weak) supernormal cones named by us "Isac's cones", through the agency of the (weak) completeness instead of compactness (see, for instance, Isac, G., 1981, 1983, 1985, 1994, 1998, Isac, G., Postolică, V., 1993, Postolică, V., 1989,1993,1994,1995, 1997, 1999, 2001, Truong, X. D. H., 1994 and so on). We use the previous bibliography and supplementary references specified in the passage below.

Theorem 1. (Bahya A. O., 1989) *A convex and normal cone K in a Hausdorff locally convex space is supernormal if and only if every net of K weakly convergent to zero converges to zero in the locally convex topology.*

Let us consider some pertinent examples

1. Any convex, closed and pointed cone in an arbitrary usual Euclidean space R^k with k in N^* is supernormal.

2. In every locally convex space any well based convex cone (i.e. generated by a convex, bounded set which does not contain the origin in its closure) is supernormal.

3. A convex cone is supernormal in a normed space if and only if it is well based.

4. Let $n \in N^*$ be arbitrary fixed and let Y be the space of all real symmetric (n, n) matrices ordered by the pointed, convex cone $C = \{A \in Y: x^T A x \geq 0, \forall x \in R^n\}$. Then, Y is a real Hilbert space with respect to the scalar product defined by $\langle A, B \rangle = \text{trace}(A \cdot B)$ for all $A, B \in Y$ and C is well based by $B = \{A \in C: \langle A, I \rangle = I\}$ where I denotes the identity matrix.

5. Every pointed, locally or weakly locally compact convex cone in any Hausdorff locally convex space is supernormal.

6. A convex cone is supernormal in a nuclear space (Pietch, A., 1972) if and only if it is a normal cone.

7. In any Hausdorff locally convex space a convex cone is weakly supernormal if and only if it is weakly normal.

8. In $L^p([a, b])$, ($p \geq 1$), the convex cone

$K_p = \{x \in L^p([a, b]) : x(t) \geq 0 \text{ almost everywhere}\}$ is supernormal if and only if $p=1$, being well based in this case by the set $B_1 = \{x \in K_1: \int_a^b x(t) dt = 1\}$. Indeed, if $p > 1$, then the sequence (x_n) defined by

$$x_n(t) = \begin{cases} n^{1/p}, & a \leq t \leq a + (b-a)/2n \\ 0, & a + (b-a)/2n < t \leq b \end{cases} \quad n \in N$$

converges to 0 in the weak topology but not in the usual norm topology. Therefore, by virtue of Theorem 1, K_p is not an Isac's cone. Generally, for every $p > 1$, K_p has as a base $B = \{x \in K_p : \int_a^b x(t) dt = 1\}$ which is unbounded and any convex cone generated by a closed and bounded set

$B_t = \{x \in B : \int_a^b |x(t)|^p dt \leq t\}$ with $t=0$ is certainly supernormal. A similar result holds for $L^p(\mathbb{R})$. Thus, if we consider a countable family (A_n) of disjoint sets which covers \mathbb{R} such that $\mu(A_n) = 1$ for all n in \mathbb{N} , where μ is the Lebesgue measure, then the sequence (y_n) given by $y_n(t) = 1$ if $t \in A_n$ and $y_n(t) = 0$ for $t \in \mathbb{R} \setminus A_n$ converges weakly to zero while it is not convergent to zero in the norm topology. Taking into account the above theorem, it follows that the usual positive cone in $K_p(\mathbb{R})$ is not a supernormal cone if $p > 1$, that is, it is not well based in all these cases. However, these cones are normal for every $p \geq 1$. The same conclusion concerning the non-supernormality is valid for the positive orthant of the usual Orlicz spaces.

9. In l^p ($p \geq 1$) equipped with the usual norm $\|\cdot\|_p$ the positive cone

$C_p = \{(x_n) \in l^p : x_n \geq 0 \text{ for all } n \in \mathbb{N}\}$ is also normal with respect to the norm topology, but it is not a supernormal cone excepting the case $p = 1$. Indeed, for every $p > 1$, the sequence (e_n) having 1 at the n th coordinate and zeros elsewhere converges to zero in the weak topology, but not in the norm topology and by virtue of Theorem 1 it follows that C_p is not supernormal. For $p = 1$, C_p is well based by the set $B = \{x \in C_1 : \|x\|_1 = 1\}$ and Proposition 5 in [32] ensures that it is supernormal. If we consider in this case the locally convex topology in l^1 defined by the seminorms $p_n((x_k)) = \sum_{k=0}^n |x_k|$ for every (x_k) in l^1 and $n \in \mathbb{N}$, which is weaker than its usual weak topology, then the usual positive cone remains supernormal with respect to this topology (now it is normal in a nuclear space and one applies Proposition 6 of [32]) but it is not well based. Taking into account the concept of H -locally convex space introduced by Precupanu, T. in 1969 and defined as any Hausdorff locally convex space with the seminorms satisfying the parallelogram law and the property that every nuclear space is also a H -locally convex space with respect to an equivalent system of seminorms [56], the above example shows that in a H -locally convex space a proper convex cone may be an Isac's cone without to be well based. Moreover, if we consider in l^2 the H -locally convex topology induced by the seminorms

$$\tilde{p}_n((x_k)) = \left(\sum_{i \geq n} |x_i|^2 \right)^{1/2}, \quad n \in \mathbb{N}, (x_k) \in \ell^2,$$

then the convex cone $C_2 = \{(x_k) \in l^2 : x_k \geq 0 \text{ for all } k \in \mathbb{N}\}$ is normal in the H -locally convex space $(l^2, \{\tilde{p}_n\}_{n \in \mathbb{N}})$, but it is not a supernormal cone because the same sequence (e_k) is weakly convergent to zero while $(\tilde{p}_n(e_k))$ is convergent to 1 for each $n \in \mathbb{N}$ and one applies again Theorem 1. Another interesting example of normal cone in a H -locally convex space which is not supernormal is the

usual positive cone in the space $L^2 \text{ loc } (R)$ of all functions from R to C which are square integrable over any finite interval of R , endowed with the system of seminorms $\{\bar{p}_n : n \in N\}$ defined by $\bar{p}_n(x) = \left(\int_{-n}^n |x(t)|^2 dt \right)^{1/2}$ for every x in $L^2 \text{ loc } (R)$. In this case, the sequence (x_k) given by:

$$x_k(t) = \begin{cases} 0, & t \in (-\infty, 0) \cup (1/k, +\infty) \\ \sqrt{k}, & t \in [0, 1/k] \end{cases}$$

converges weakly to zero, but it is not convergent in the H -locally convex topology. The results follows by Theorem 1. It is clear that every weak topology is a H -locally convex topology and, in these cases, the supernormality of convex cones coincides with the normality thanks to the Corollary of Proposition 2 in [32].

10. In the space $C([a, b])$ of all continuous, real valued functions defined on every non-trivial, compact interval $[a, b]$ equipped with the usual supremum norm the convex cone $K = \{x \in C([a, b]) : x \text{ is concave, } x(a)=x(b) = 0 \text{ and } x(t) \geq 0 \text{ for all } t \in [a, b]\}$ is supernormal, being well based by the set $\{x \in K : x(t_0) = 1\}$ for some arbitrary $t_0 \in [a, b]$. The hypothesis that all $x \in K$ are concave is essential for the supernormality.

11. The convex cone of all nonnegative sequences in the space of all absolutely convergent sequences is the dual of the usual positive cone in the space of all convergent sequences. Consequently, it has a weak star compact base and hence it is a weak star supernormal cone.

12. In l^∞ or in c_0 equipped with the supremum norm, the convex cone consisting of all sequences having all partial sums non-negative is not normal, hence it is not supernormal.

13. In every Hausdorff locally convex space any normal cone is supernormal with respect to the weak topology.

14. In every locally convex lattice which is a (L) -space the ordering cone is supernormal (see also the Example 7 given by Isac, G. in 1994).

15. If we consider the space of all locally integrable functions on a locally compact space Y with respect to a Radon measure μ endowed with the topology induced by the family of seminorms $\{p_A\}$ where $p_A(f) = \int_A |f(x)| d\mu$ for every non-empty and compact subset A of Y and every locally integrable function f , then the convex cone $K = \{f : f(x) \geq 0, \forall x \in Y\}$ is supernormal.

16. If Z is any locally convex lattice ordered by an arbitrary convex cone K and Z^* is its topological dual ordered by the corresponding dual cone K^* , then the cone K is supernormal with respect to the locally convex topology defined on Z by the neighbourhood base at the origin $\{[-f, f]^0\}_{f \in \bar{K}}$

17 In every regular vector space (E, K) (that is, the order dual E^* separates the points of E) with the property that $E = K - K$ the convex cone K is supernormal with respect to the topology defined in the preceding example.

18. Any *semicomplete cone* in a Hausdorff locally convex space is *supernormal* (for this concept see the Example 11 of [35]).

Remark 1. Clearly, if a convex cone K is *supernormal* in a normed space, then K admits a strictly positive, linear and continuous functional, that is, there exists a linear, continuous functional f such that $f(k) > 0$ for all $k \in K \setminus \{0\}$. Generally, the converse is not true even in a Banach space as we can see in the following examples:

19. If one considers in the usual space l^p ($1 \leq p \leq \infty$) the convex cone $K = l^p_+ = \{x = (x_i) \in l^p : x_i \geq 0 \text{ for every } i \in N\}$ of infinite vectors with non-negative components, then the functional φ defined by $\varphi(k) = \sum_{i=1}^{\infty} k_i$ for any $k = (k_i) \in l^p$ is linear, continuous and strictly positive. But, as we have seen in the above considerations (Example 9), this cone is *supernormal* if and only if $p = 1$.

20. Let K be the usual positive cone $L^p_+ = \{x \in L^p([a, b]) : x(t) \geq 0 \text{ almost everywhere}\}$ in $L^p([a, b])$ ($1 \leq p \leq \infty$). Then, the linear and continuous functional ψ on $L^p([a, b])$ given by $\psi(x) = \int_a^b x(t)dt$ for every $x \in L^p([a, b])$ is strictly positive on K while K is *supernormal* (see the above Example 8) if and only if $p = 1$.

Therefore, l^1_+ and L^1_+ are *supernormal cones* with empty topological interiors and for every $p \in (1, +\infty)$ it follows that l^p_+ and L^p_+ are *normal cones* with empty interiors which are not *supernormal*. Hence, these convex cones are not well based. A very simple example of *supernormal cones* having non-empty topological interior is R^n_+ ($n \in N^*$).

Remark 2. In the order complete vector lattice $B([a, b])$ of all bounded, real valued functions on a compact non-singleton interval $[a, b]$ endowed with its usual norm the standard positive cone $K = \{u \in B([a, b]) : u(t) \geq 0 \text{ for all } t \in ([a, b])\}$ is *normal* but it has not a base, that is, it is not *supernormal*. However, this cone has non-empty interior. If we consider the linear space l^l endowed with the separated locally convex topology generated by the family $\{p_n : n \in N\}$ of seminorms defined by

$$p_n(x) = \sum_{k=0}^n |x_k|$$

for every $x = (x_k) \in l^l$, then the convex cone

$K = \{x = (x_k) \in l^l : x_k \geq 0 \text{ whenever } k \in N\}$ is *supernormal* but it is not well based.

Remark 3. The natural context of *supernormality* (nuclearity) for convex cones is any separated locally convex space. Isac, G. introduced the concept of “nuclear cone” in 1981, published it in 1983 and he showed that in a normed space a convex cone is *nuclear* if and only if it is well based or equivalently iff it is “with plastering”, the last concept being defined by Krasnoselski, M. A. in fifties (see, for example, Krasnoselski, M. A., 1964 and so on). Such a convex cone was initially called “nuclear cone” by Isac, G. (1981) because in every nuclear space [56] any normal

cone is a nuclear cone in Isac's sense (Proposition 6 of [32]). Afterwards, since the nuclear cone introduced by Isac appears as a reinforcement of the normal cone, it was called *supernormal*. The class of supernormal cones in Hausdorff locally convex spaces was initially imposed by the theory and the applications of the efficient (Pareto minimum type) points (especially existence conditions based on completeness instead of compactness were decisive together with the main properties of the efficient points sets), the study of critical points for dynamical systems and conical support points and their importance was very well illustrated by important results, examples and comments in the specified references and in other connected papers. It is also very significant to mention again that the concept of supernormality introduced by , [31] is not a simple generalization of the corresponding notion defined in normed linear spaces by Krasnoselski, M. A. and his colleagues in the fifties. Thus, for example, Isac's supernormality attached to the convex cones has his sense in every Hausdorff locally convex space identically with the well known Grothendieck's nuclearity. By analogy with the fact that a normed space is nuclear in Grothendieck's sense if and only if it is isomorphic with an usual Euclidean space, a convex cone is supernormal in a normed space if and only if it is well based, that is, it is generated by a convex bounded set which does not contain the origin in its closure. Therefore, the more appropriate background for the supernormal cones is any separated locally convex space.

6. USEFUL SPLINES FOR THE BEST APPROXIMATION AND OPTIMIZATION IN H-LOCALLY CONVEX SPACES

We conclude this research report with some topics on the best approximation (simultaneous and vectorial) and the optimization in H -locally convex spaces. So, it is known that the concept of H -locally convex space was introduced and studied for the first time by [73] and defined as any Hausdorff locally convex space with the seminorms satisfying the parallelogram law. At the same time, we introduced the notion of spline function in H -locally convex space [57] and we established the basic properties of approximation and optimal interpolation for these splines. Our splines are natural extensions in H -locally convex spaces of the usual abstract splines which appear in any Hilbert space like the minimizing elements for a seminorm subject to the restrictions given by a set of linear continuous functionals. Let $(X, P = \{p_\alpha : \alpha \in I\})$ be a H -locally convex space with each seminorm p_α being induced by a scalar semiproduct $(\cdot, \cdot)_\alpha (\alpha \in I)$ and M a closed linear subspace of X for which there exist a H -locally convex space $(Y, Q = \{q_a : a \in I\})$ with each seminorm $q_a \in Q$ generated by a scalar semiproduct $\langle \cdot, \cdot \rangle_a (\alpha \in I)$ and a linear (continuous) operator $U : X \rightarrow Y$ such that $M = \{x \in X : (x, y)_\alpha = \langle Ux, Uy \rangle_a, \forall a \in I\}$.

The space of spline functions with respect to U was defined by Postolică, V.(1981) as the U -orthogonal of M , that is,

$M^\perp = \{x \in X : \langle U_x, U_\zeta \rangle_\alpha = 0, \forall \zeta \in M, \alpha \in I\}$. Clearly, M^\perp is the orthogonal of M in the H -locally convex sense.

Let $x_0 \in X$ and G a non-empty subset of X .

Definition 1. [34] $g_0 \in G$ is said to be a best simultaneous approximation for x_0 by the elements of G with respect the family P (abbreviated g_0 is a P -b.s.a. of x_0) when $p_\alpha(x_0 - g_0) \leq p_\alpha(x_0 - g)$ for all $g \in G$ and $p_\alpha \in P$.

If, in addition, each element $x \in X$ possesses at least one P -b.s.a. in G , then the set G is called P -simultaneous proximal.

Definition 2. [34] $g_0 \in G$ is said to be a best vectorial approximation of x_0 by G with respect to P (abbreviated g_0 is a P -b.v.a. of x_0) if $(p_\alpha(x_0 - g_0)) \in \text{MIN}_K(\{(p_\alpha(x_0 - g)) : g \in G\})$ where $K = \mathbb{R}_+^1$.

If, in addition, each element $x \in X$ possesses at least one P -b.v.a. in G , then the set G is called P -vectorial proximal.

Let us consider the direct sum $X' = M \oplus M^\perp$ and for every $x \in X'$, we denote its projection onto M^\perp by s_x . Then, taking into account the Theorem 4 obtained by Postolică, V. in 1981, it follows that this spline is a best simultaneous U -approximation of x with respect to M^\perp since it satisfies all the next conditions : $p_\alpha(x - s_x) \leq p_\alpha(x - y) \forall y \in M^\perp, p_\alpha \in P$.

Moreover, following the definition of the approximate efficiency, the results given in Chapter 3 of (Isac, G., Postolică, V., 1993) and the conclusions obtained by (Postolică, V., 1981, 1993, 1998), we have

Theorem 1.

(i) for every $x \in X$ the only elements of best simultaneous and vectorial approximation with respect to any family of seminorms which generates the H -locally convex topology on X by the linear subspace of splines are the spline functions s_x . Moreover, if M and M^\perp supply an orthogonal decomposition for X , that is $X = M \oplus M^\perp$, then M^\perp is simultaneous and vectorial proximal;

(ii) if $K = \mathbb{R}_+^1$, then for each $s \in M^\perp$, every $\sigma \in M^\perp$ is the only solution of following optimization problem $\text{MIN}_K(\{(q_\alpha(U(\eta - s))) : \eta \in X' \text{ and } \eta - \sigma \in M\})$;

(iii) for every $x \in X'$ its spline function s_x is the only solution for the next vectorial optimization problems:

$$\text{MIN}_K(\{(q_\alpha(U(\eta - x))) : \eta \in M^\perp\}), \text{MIN}_K(\{(p_\alpha(x - y)) : y \in M^\perp\}), \text{MIN}_K(\{(q_\alpha(Uy)) : y - x \in M\}).$$

Finally, let us consider two numerical examples in which, following

[34], [57], [64] we specify the expressions of splines and M and M^\perp realize orthogonal decompositions.

Example 1. Let $X = H^m(\mathbb{R}) = \{f \in C^{m-1}(\mathbb{R}) : f^{m-1} \text{ is locally absolutely continuous and } f^m \in L^2_{loc}(\mathbb{R})\}, m \geq 1$ endowed with the H -locally convex topology generated by the scalar semiproducts

$$(x, y)_k = \sum_{h=0}^{m-1} [x^h(k)y^h(k) + x^h(-k)y^h(-k)] + \int_{-k}^k x^m(t)y^m(t)dt, k = 0, 1, 2, \dots \text{ and}$$

$Y = L^2_{loc}(R)$ with the H-locally convex topology induced by the scalar semiproducts $\langle x, y \rangle_k = \int_{-k}^k x(t) y(t) dt, k=0,1,2, \dots$

If $U: X \rightarrow Y$ is the derivation operator of order m , then

$$M = \{x \in H^m(R) : x^{(h)}(n) = 0, \forall h = \overline{0, m-1}, n \in Z\}$$

$$\text{and } M^\perp = \{s \in H^m(R) : \int_{-k}^k s^{(m)}(t) x^{(m)}(t) dt, \forall x \in M, k=0,1,2, \dots\}$$

We proved in (Postolică, V., 1981) that

$M^\perp = \{s \in H^m(R) : s(n, n+1)$ is a polynomial function of degree $2m-1$ at most} and if $y = (y_v), y' = (y'_v), y'' = (y''_v), y^{(m-1)} = (y^{(m-1)}_v)$ are m sequences of real numbers, then there exists a unique spline $S \in M^\perp$ satisfying the following conditions of interpolation: $S^{(h)}(n) = y^{(h)}(n)$ whenever $h = \overline{0, m-1}$ and $v \in Z$. Moreover, we observed in the paragraph 3 of (Isac, G., Postolică, V., 1993) that any spline function S such as this is defined by

$$S(x) = p(x) + \sum_{h=0}^{m-1} c_1^h(x-1) + 2m-1 + \sum_{h=0}^{m-1} c_2^h(x-2) + 2m-1 + \dots + \sum_{h=0}^{m-1} c_0^h(-x) + 2m-1 + \dots$$

where $u+ = (|u|+u)/2$ for every real number u , p is a polynomial function of degree $2m-1$ at most perfectly determined by the conditions $p^{(h)}(0) = y_0^h$ and $p^{(h)}(1) = y_1^h$ for all $h = \overline{0, m-1}$ and the coefficients $c_v^h (h = \overline{0, m-1}, v \in Z)$ are successively given by the general interpolation.

Therefore, for every function $f \in H^m(R)$, there exists a unique function denoted by $S_f \in M^\perp$ such that $S_f^{(h)}(n) = f^{(h)}(n), \forall h = \overline{0, m-1}$ and $n \in Z$. Hence, in this case, M and M^\perp give an orthogonal decomposition for the space $H^m(R)$.

Example 2. Let $X = F_m = \{f \in C^{m-1}(R) : f^{(m-1)}$ is locally absolutely continuous and $f^{(m)} \in L^2(R)\}$ endowed with the H-locally convex topology induced by the scalar semiproducts

$(x, y)_v = x(v)y(v) + \int_R x^{(m)}(t)y^{(m)}(t)dt, v \in Z, Y = L^2(R)$ with the topology generated by the inner product $(x, y)_n = \int_R x(t) y(t)dt, n \in Z$ and $U: X \rightarrow Y$ be the derivation operator of order m . Then, $M = \{x \in F_m : x(n) = 0 \text{ for all } n \in Z\}$ and

$$M^\perp = \{s \in F_m : \int_R x^{(m)}(t) y^{(m)}(t) dt = 0 \text{ for every } x \in M\}.$$

In a similar manner as in Example 1 it may be proved that M^\perp coincides with the class of all piecewise polynomial functions of order $2m$ (degree $2m-1$ at most) having their knots at the integer points. Moreover, for every function f in F_m there exists a unique spline function $S_f \in M^\perp$ which interpolates f on the set Z of all integer numbers, that is, S_f satisfies the equalities $S_f(n) = f(n)$ for every $n \in Z$, being defined by

$$S_f(x) = p(x) + a_1(x-1) + 2m-1 + a_2(x-2) + 2m-1 + \dots + a_0(-x) + 2m-1 + a_{-1}(-x-1) + 2m-1 + \dots$$

where $u+$ has the same signification as in Example 1, the coefficients $a_v (v \in Z)$ are successively and completely determined by the interpolation conditions $S_f(v) =$

$f(v), v \in \mathbb{Z} \setminus \{0, 1\}$ and p is a polynomial function satisfying the conditions $p(0)=f(0)$ and $p(1)=f(1)$. The uniqueness of S_f is ensured in Theorem 2 given by [57].

Thus, M and M^\perp give an orthogonal decomposition of the space F_m and, as in the preceding example, M^\perp is simultaneous and vectorial proximal with respect to the family of seminorms generated by the above scalar semiproducts.

Remark 1. Our examples show that the abstract construction of splines can be used to solve also several frequent problems of interpolation and approximation, having the possibility to choose the spaces and the scalar semiproducts. It is obvious that for a given (closed) linear subspace of a H -locally convex space X such a H -locally convex space Y (respectively, a linear (continuous) operator $U: X \rightarrow Y$) would not exist. Otherwise, the problem of best vectorial approximation by the corresponding orthogonal space of any (closed) linear subspace M for the elements in the direct sum $M \oplus M^\perp$ might be always reduced to the best simultaneous approximation. But, in general, such a possibility doesn't exist. Even in a H -locally convex space it is possible that there exist best vectorial approximations and the set of all best simultaneous approximations to be empty for some element of the space. We confine ourselves to mention the following simple example.

Example 3. Let $X = \mathbb{R}^N$ endowed with the topology generated by the family $P = \{p_i : i \in \mathbb{N}\}$ of seminorms defined by $p_i(x) = |x_i| (i \in \mathbb{N})$ for every

$x = (x_i) \in X$ and $G = \{(x_i) \in X : x_i \geq 0 \text{ whenever } i \in \mathbb{N} \text{ and } \sum_{i \in \mathbb{N}} x_i = 1\}$

X is a P -simultaneous strictly convex (Isac, G., Postolici-ÅL, V., 1993)

H -locally convex space. Nevertheless, every element of G is P -b.v.a. for the origin while its corresponding set of the best simultaneous approximations with respect to P is empty.

References

- [1] ARROW, K.J. *Social choice and individual values*, The second edition. Yale University Press, (1963).
- [2] ARROW, K.J., HAHN, F.M. – *General competitive analysis*, Holden Day, place State New York, (1971).
- [3] AUBIN, J.P. – *Optima and equilibria. An introduction to nonlinear analysis*, Springer – Verlag Berlin Heidelberg, (1993).
- [4] BAHYA, A. O. - *Ensembles Coniquement Bornés et CĂnes Nucléaires dans les Éspaces Localement Convexes Separés*. Thésés de Doctorat de 3-ème Cycle en Mathématiques. L'École Normale Supérieure - Takaddoum Rabat (Maroc), (1989).
- [5] BENSON, H.P. – *The vector maximization problem: proper efficiency and stability*, SIAM J. Appl. Math. 32, (1977), 64-72.
- [6] BENSON, H.P. – *An improved definition of proper efficiency for vector minimization with respect to cones*, J. Math. Anal. Appl. 79, (1979), 232-241.
- [7] BENSON, H.P. – *Efficiency and proper efficiency for vector minimization with respect to cones*, J. Math. Anal. Appl. 93, (1983), 273-289.
- [8] BOBOC, N., BUCUR, GH. - *Convex cones of continuous functions on compact spaces*, (in Romanian), Publishing House of Romanian Academy, Bucharest, (1976).

- [9] BORWEIN, J.M. – *Proper efficient points for maximization with respect to cones*, SIAM J. Control Optim., 15, (1977), 57 – 63.
- [10] BORWEIN, J.M. – *On the existence of Pareto efficient points*, Math Oper. Res. 8, no. 1, (1983), 64 - 73.
- [11] BRUNDTLAND COMMISSION (WORLD COMMISSION ON ENVIRONMENT AND DEVELOPMENT) – *Our common future*, Oxford University Press, Oxford, (1987).
- [12] BUCUR, I., POSTOLICĂ, V.- *A coincidence result between sets of efficient points and Choquet boundaries in separated locally convex spaces*, Research Report at the 4th Workshop of the German Working Group on Decision Theory, Hotel Talblick, Holzgau, country-region Germany, March 14-18, 1994. Optimization, 36, (1996), 231-234.
- [13] CARDINALI, T. and PAPALINI, F.- *Fixed point theorems for multifunctions in topological vector spaces*, J. Math. Anal. Appl., 186, (1994), 769-777.
- [14] CHOQUET, G.- *Ensembles et \mathcal{A} nes convexes faiblement complets*, C. R. Acad. Sci. Paris, 254, (1962), 2123-2125.
- [15] CHOQUET, G.- *Theory of capacities*, Ann. Inst. Fourier, 5, (1955).
- [16] CHOQUET, G., DENY, J.- *Ensembles semi-reticulés et ensembles reticulés des fonctions continues*, Journ. Math. Pures Appl., 36, (1957), 179-189.
- [17] CHOQUET, G., MEYER, P. A.- *Existence et unicité des représentations dans les convexes compacts quelconques*. Ann. Inst. Fourier, 13, (1963), 139-154.
- [18] DAUER, J. P., GALLAGHER, R. J. – *Positive proper efficient points and related coneresults in vector optimization theory*, SIAM J. Control Optim. 28(1), (1990), 158-172.
- [19] DEBREU GERARD – *Theory of value*, Wiley, New York, (1959).
- [20] DOMINIAK CEZARY – *Multicriteria decision aid under uncertainty*, In: Multiple Criteria Decision Making'05 (Ed.Tadeusz Trzaskalik), Publisher of The Karol Adamiecki PlaceTypeUniversity of Economics in Katowice, (2006), 63-81.
- [21] EHRGOTT, MATTHIAS – *Multicriteria Optimization*, Springer Berlin-Heidelberg, (2005).
- [22] ENGAU, StateAL. – *Definition and Characterization of Geoffrion Proper Efficiency for Real Vector Optimization with Infinitely Many Criteria*, J. Optim. Theory Appl (2015) 165: 439 – 457.
- [23] FISHBURN, P. C. – *Foundations of risk management science placeI. Risk or probable loss*. Management Science, 30, (1984), 396 – 406.
- [24] GASS SAUL, placeI., HARRIS CARL, M. – *Enciclopedia of operations research and management science*, Centennial Edition. Kluwer Academic Publishers, Boston, Dordrecht, London, (2001).
- [25] GEOFFRION, A. M.- *Proper efficiency and the theory of vector maximzation*, J. Math. Anal. Appl., 22, (1968), 618 -630.
- [26] HANSSON, S.O. – *Welfare, justice and Pareto efficiency*, Ethical Theory and Moral Practise no. 7, (2004).
- [27] HARTLEY, R. – *On cone - efficiency, cone convexity and cone – compactness*, SIAM J. Appl. Math., 34(2), (1978), 211-222.
- [28] HENING, M. I. – *Proper efficiency with respect to cones*, J. Optim. Theory. Appl., 36, (1982), 387-407.
- [29] HEYNE, PAUL – *Efficiency*, Research Report, University of Washington, (2000).
- [30] HYERS, D. H., ISAC, G., RASSIAS, T. M. – *Topics in nonlinear analysis and applications*, World Scientific Publishing Co, Pte. Ltd., (1997).

- [31] ISAC, G. - *Points critiques des systĂŠmes dynamiques. CĂŞnes nuclĂŞsaires et optimum de Pareto*, Research Report, Royal Military College of St. Jean, QuĂŞbec, Canada, (1981).
- [32] ISAC, G. - *Sur l'existence de l'optimum de Pareto*, Riv. Mat. Univ. Parma, 4(2),(1983), 303 - 325.
- [33] ISAC, G. - *Supernormal cones and absolute summability*, Libertas Mathematica, 5, (1985), 17 - 32.
- [34] ISAC, G., POSTOLICĂ, V.- *The best approximation and optimization in locally convex spaces*, Verlag Peter Lang GmbH, Frankfurt am Main, Germany, (1993).
- [35] ISAC, G.- *Pareto optimization in infinite dimensional spaces: the Importance of nuclear cones*, J. Math. Anal. Appl., 182(A201), (1994), 393 - 404.
- [36] ISAC, G. - *On Pareto efficiency. A general constructive existence principle*, Research Report (16 pp.), Department of Mathematics and Computer Science, Royal Military College of Canada, (1998).
- [37] ISAC, G., BAHYA, A., O. - *Full nuclear cones associated to a normal cone. Application to Pareto efficiency*, Applied Mathematics Letters, 15, (2002), 633 - 639.
- [38] ISAC, G., BULAVSKY, V. A., KALASHNIKOV, V. V.- *Complementarity, equilibrium, efficiency and economics*, Kluwer Academic Publishers, (2002).
- [39] ISAC, G., POSTOLICĂ, V.- *Full nuclear cones and a relation between strong optimization and Pareto efficiency*, J. Global Optimiz., 32, (2005),507 - 516.
- [40] JABLONSKY, J.- *A slack based model for measuring super-efficiency in data envelopment analysis*, In: Multiple Criteria Decision Making'05 (Ed.Tadeusz Trzaskalik), Publisher of The Karol Adamiecki University of Economics in Katowice, (2006), 101-112.
- [41] KORHONEN,P., LUPTĂÈIK, M. - *Eco-efficiency analysis of power plants : an extension of data envelopment analysis*, European Journal of Operational Research, 154, (2004), 437 - 446.
- [42] KOSMIDOU, K., ZOPOUNIDIS, C. - *Managing interest rate risk in commercial banks via multicriteria analysis*, Research Paper presented at The 61st Meeting of The European Working Group "Multiple Criteria Decision Aiding", University of Luxembourg, March 10 - 11, (2005).
- [43] KRAMAR, E. - *Locally convex topological vector spaces with hilbertian seminorms*, Rev. Roum. Math Pures et Appl., Tome XXVI, no. 1, (1981), 55-62
- [44] KRASNOSELSKI, M. A. - *Positive Solutions of Operator Equations*. GrĂŞnningen, Noordhoff, (19640).
- [45] KUHN, H. W., TUCKER, A. W. - *Nonlinear programming*. Proceedings of the Second Berkely Sympsiom on Mathematical Statistics and Probability (Neyman J. Ed.), University of California Press, Berkely, (1950), 481-492.
- [46] LORIDAN, P. - *ϵ -solutions of vector minimization problems*, J. Optim. Theory Appl., 43(2), (1984), 265 - 276.
- [47] LUC, D. T. - *Theory of vector optimization*, Springer-Verlag, (1989).
- [48] LUPTĂÈIK, M., BÖHM, B. - *Measuring eco - efficiency an a Leontief input - output model*, In: Multiple Criteria Decision Making'05 (Ed.Tadeusz Trzaskalik), Publisher of The Karol Adamiecki University of Economics in Katowice, (2006), 121-135.
- [49] NACHBIN, L. - *Topology and order*, StateplacNew York, Van Nostrand, (1965).
- [50] NÉMETH, A. B. - *Between Pareto efficiency and Pareto ϵ -efficiency*, Optimization,10(5) , (1989), 615 - 637.
- [51] NG., K. F., ZHENG, X. Y. - *Existence of efficient points in vector optimization and generalized Bishop-Phelps theorem*, JOTA, 115, (2002), 29 - 47.

- [52] NYBERG, L. et al – *Förslag till Transportmål för Stockholms Län (Suggested Transport Goals for Stockholms PlaceTypeCounty)*, Stockholm: Länsstyrelsen i Stockholms Län, (2001).
- [53] PARK, S. – *Some coincidence theorems on acyclic multifunctions applications*. In: Tan, K.-K (ed) *Fixed Point Theory and Applications*, World Sci. NJ, (1992), 248–277.
- [54] PARETO, VILFREDO – *Manuel d'économie politique*, Girard et Brière, (1909).
- [55] PERESSINI, A. L. - *Ordered topological vector spaces*, Harper 8 Row, New York State, (1967).
- [56] PIETCH, A. – *Nuclear Locally Convex Spaces*. Springer -Verlag, (1972).
- [57] POSTOLICĂ, V. - *Spline functions in H-locally convex spaces*, An. Șt. Univ. "Al. I. Cuza", Iași, 27, (1981), 333-338.
- [58] POSTOLICĂ, V. - *Existence Conditions of Efficient Points for Multifunctions with Values in Locally Convex Spaces*. Stud. Cerc. Mat., Tom 41, no. 4, (1989), p. 325-331.
- [59] POSTOLICĂ, V. - *New existence results for efficient points in locally convex spaces ordered by supernormal cones*, J. Global Optimiz., 3, (1993), 233-242
- [60] POSTOLICĂ, V. - *An Extension to Sets of Supernormal Cones and Generalized Subdifferentials*. Optimization, 29, (1994), p. 131-139.
- [61] POSTOLICĂ, V. - *Properties of Pareto sets in locally convex spaces*, Optimization, 34, (1995), 223-229.
- [62] POSTOLICĂ, V. - *Recent Conditions for Pareto Efficiency in Locally Convex Spaces*. Research Report at the Joint International Meeting Euro XV-INFORMS XXXIV, Barcelona, country-regionSpain, July 14-17, (1997).
- [63] POSTOLICĂ, V. - *Properties of efficient points sets and related topics*, Research Report at The Second International Conference on Multi-Objective Programming and Goal Programming, Torremolinos, country-regionSpain, May 16-18, 1996. Published in *Advances in Multiple Objective and Goal Programming, Lecture Notes in Economics and Mathematical Systems*, 455, (Eds.: Rafael Caballero, Francisco Ruiz, Ralph E. Steuer), (1997), 201-209.
- [64] POSTOLICĂ, V. - *A method which generates splines in H-locally convex spaces and connections with vectorial optimization*, Positivity, 2 (4), (1998), 369 - 377.
- [65] POSTOLICĂ, V. - *Conditions for Pareto efficiency in locally convex spaces*, Mathematical Reports of Romanian Academy 2, 1(51), April-June, (1999), 257-269.
- [66] POSTOLICĂ, V., SCARELLI, A. - *Some connections between best approximation, vectorial optimization and multicriterial analysis*, Nonlinear Analysis Forum, 5, (2000), 111-123.
- [67] POSTOLICĂ, V. SCARELLI, A., VENZI, L. - *On the equilibrium of a multidimensional ecosystem*, Nonlinear Analysis Forum, 6(2), (2001), 321-335.
- [68] POSTOLICĂ, V. - *New Existence Results of Solutions for Vector Optimization Programs with Multifunctions*. Optimization, vol. 50, (2001), p. 253-263.
- [69] POSTOLICĂ, V. - *Pareto efficiency, Choquet boundaries and operators in Hausdorff locally convex spaces*, Nonlinear Analysis Forum, 7(2), (2002), 215 – 230.
- [70] POSTOLICĂ, V. – *A class of generalized dynamical systems in connection with Pareto efficiency and related topics*, Research Paper submitted to the International Conference "Complex Systems, Intelligence and Modern Technological Applications", Cherbourg, country-regionFrance, September 19 - 22, (2004). Published in the corresponding Electronics Proceedings Volume.
- [71] POSTOLICĂ, V. - *Choquet boundaries and efficiency*, Computer and Mathematics with Applications, 55, (2008), 381 - 391.
- [72] POSTOLICĂ, V. – *Efficiency and recent applications*. Research Report and Poster at XVI International Congress on Mathematical Physics, Prague, August 3 – 8, (2009).

- [73] PRECUPANU, T. - *Espaces linéaires à semi-normes hilbertiennes*, An. Șt. Univ. "Al. I. Cuza", Iași, 15, (1969), 83-93.
- [74] PRECUPANU, T. - *Scalar minimax properties in vectorial optimization*, International Series of Numerical Mathematics, Birkhäuser Verlag Basel, 107, (1992), 299 - 306.
- [75] ROSENCRANTZ, HOLGER – *Transport policy goals, welfare concerns, and rationality*, Research Report presented at The 62-nd Meeting of The European Working Group "Multiple Criteria Decision Aiding", Borlänge, Sweden, 22-24 September, (2005).
- [76] ROTMANS, JAN – *New trends in modelling: complexity, sustainable development, ethics*, The International Centre for Integrative Studies (ICIS). Plenary Talk at the 12th Mini Euro Conference organized by the Euro Working Groups Decision Support Systems (DSS), Human Centered Processes (HCP) and Prometheus (Ethics), Vrije University of Brussels, Belgium, April 2-5, (2002).
- [77] SONGSAK SRIBOONCHITTA, WING-KEUNG WONG, SOMPONG DHOMPONGSA, HUNG, T. NGUYEN – *Stochastic dominance and applications to finance, risk and economics*. CRC Press, Taylor and Francis Group, October (2009), 456 pp., ISBN: 978-1-4200-8266-1.
- [78] SONNTAG, Z., ZĂLINESCU, C. - *Comparison of existence results for efficient points*, JOTA, 105, (2000), 161-188.
- [79] STERNA - KARWAT, A. - *On the existence of cone maximal points in real topological linear spaces*, Israel Journal of Mathematics, 54(1), (1986), 33 - 41.
- [80] STEWART, D. J. – *Uncertainties in MCDA*, In: Multi Criteria Decision Analysis (Ed. P. Greco). Springer-Verlag, (2004).
- [81] TAYLOR, A. D. – *Mathematics and politics. Strategy, voting, power and proof*. Springer – Verlag New York, Inc., (1995).
- [82] TRUONG, X. D. H. - *A note on a class of cones ensuring the existence of efficient points in bounded complete sets*, Optimization, 31, (1994), 141-152.
- [83] TRUONG, X. D. H. - *On the existence of efficient points in locally convex spaces*, J. Global Optimiz., 4, (1994), 265-278.
- [84] www.wikipedia.org
- [85] ZHANG CONG - JUN – *Generalized bi-quasi-variational inequalities and generalized quasi-variational inequalities*, Acta Analysis Functionalis Applicata, 7(2), (2005), 116 - 122 .
- [86] ZIMMERMANN, H. – *An application - oriented view of modeling uncertainty*, European Journal of Operation Research, 122, (2000), 190 - 198.